

ORGANISATION DE L'UNITE  
AFRICAINNE

—•••—

COMMISSION SCIENTIFIQUE  
TECHNIQUE ET DE LA RECHERCHE

—•••—

CONSEIL PHYTOSANITAIRE  
INTERAFRICAIN

—



ORGANIZATION OF AFRICA  
UNITY


—•••—

SCIENTIFIC, TECHNICAL AND  
RESEARCH COMMISSION

—•••—

INTERAFRICAN  
PHYTOSANITARY  
COUNCIL

—



# AFRICAN JOURNAL OF PLANT PROTECTION



**REVUE AFRICAINE DE LA  
PROTECTION DES  
PLANTES**



VOLUME N° 5 (2)

Août : 1995  
August :



Edited by D M Masaba<sup>1</sup> and N Nkouka<sup>2</sup>

<sup>1</sup>Coffee Research Foundation,  
P O Box 4, RUIRU, Kenya

<sup>2</sup>InterAfrican Phytosanitary Council (OAU/IAPSC)  
P O Box 4170, YAOUNDE, Cameroon

**INTERAFRICAN PHYTOSANITARY COUNCIL  
OF THE ORGANIZATION OF AFRICAN UNITY  
(OAU/IAPSC)**

**COFFEE RESEARCH FOUNDATION  
(CRF)**

# CONTENTS

FOREWORD .....	iv
----------------	----

## WELCOME ADDRESSES:

Hon. S. Nyachae .....	vi
-----------------------	----

Dr. N. Nkouka .....	viii
---------------------	------

RESOLUTIONS AND RECOMMENDATIONS .....	x
---------------------------------------	---

## COUNTRY REPORTS:

Coffee Berry Disease in Kenya D. M. Masaba and others .....	1
--	---

Research system on Coffee Berry Disease and its control in Tanzania F. S. Ngulu .....	15
--	----

Review of current status of Coffee Berry Disease (CBD) research development in Ethiopia E. Derso .....	21
---	----

Coffee Berry Disease in Zimbabwe P. Zvoutete .....	27
---	----

Coffee Berry Disease in Malawi N. A. Phiri .....	31
---	----

Coffee Berry Disease in Rwanda André Sibomana .....	35
--	----

Coffee Berry Disease in Cameroon J. Bakala .....	41
---	----

LIST OF PARTICIPANTS .....	47
----------------------------	----

## FOREWORD

The Coffee Research Foundation (CRF), Ruiru and the Inter-African Phytosanitary Council (IAPSC) of the Organization of African Unity (OAU) jointly organized a meeting to constitute a network on research on the control of the Coffee Anthracnose, commonly known as Coffee Berry Disease (CBD). The meeting was recommended by a joint Permanent Secretariat set up by resolutions CM/RES 1173 (xlviii) and CM/RES 1238 (1) of the OAU Council of Ministers in their 48th and 58th Ordinary Sessions. Participants in the meeting were expected to come from about ten African countries which grow arabica coffee and which are affected by CBD; namely Kenya, Burundi, Rwanda, Uganda, Zimbabwe, Tanzania, Ethiopia, Malawi and Zambia; as well as representatives of the General Secretariat and the Scientific, Technical and Research Commission of the OAU. The meeting was held at Hotel 680, Nairobi between 26 and 28 May 1993.

We are all aware that Agriculture is the mainstay of the economy of most African countries. It is also a way of life for the majority of the peoples, providing employment to a high percent of the population. For many of the African countries, coffee is one of the principal export commodities. Coffee was contributing, on average, 26% of the total domestic export value of Kenya until 1988/89, when this per cent started to drop as a result of the low world prices experienced after the collapse of the price support mechanism of the International Coffee Agreement (ICA). There has also been a decline in overall production of coffee linked to the now unstable and depressed market situation resulting in low prices already alluded to above. The low prices led to poor husbandry practices and generally low morale on the part of the farmers.

The current cost of producing coffee in Kenya is about Kf 2,700 per metric tonne which is considered high. Control of the diseases alone contributes 30% of this total cost. Coffee berry disease (CBD) caused by a fungus *Colletotrichum kahawae* Waller & Bridge, is the most severe disease of coffee in Kenya and other African countries which grow Arabica coffee. The disease attacks berries in all stages of growth, but the most susceptible stages are the green expanding berries (6-16 weeks after flowering) and ripe berries, and especially when these stages coincide with wet weather conditions.

On green berries, symptoms first appear as small dark, sunken patches (anthracnose) which spread rapidly and may cover the whole berry. Usually the fungus penetrates the interior and destroys the beans. Eventually, the whole berry dries out and takes the appearance of black mummified bodies resembling 'buni'. Infected berries may also be shed off as soon as lesions develop.

Coffee berry disease was first detected in 1922 in Western Kenya, but has since spread to all coffee growing areas in Kenya. It has also been reported in many other African countries, namely, Uganda, Tanzania, Rwanda, Burundi, Zaire, Angola, Ethiopia, Cameroon, Zimbabwe, Zambia and Malawi. The disease has not been reported outside the African continent.

Rainfall is the most important facet of weather which influences CBD, because the causal agent of CBD requires water for dispersal of its spores, subsequent germination and penetration of healthy berries, leading to development of lesions. The fungus is also favoured by cool temperatures in the range of 15 to 25°C. These two weather requirements, coupled with the almost continuous flowering of coffee trees, make the high altitude coffee more vulnerable to CBD than the coffee from medium and low altitude zones.

The Coffee Research Foundation, Kenya, has made a lot of progress in developing efficient chemical control measures to CBD. These include spraying with a number of recommended fungicides to protect the developing crop during the rainy seasons. Copper-based fungicides were used in Kenya until 1969 when captafol and later other organic fungicides, became available. The small-scale farmers in Kenya have continued using copper mainly because of its relatively low cost, and also because copper controls other major coffee diseases, namely coffee leaf rust, and in some areas, bacterial blight of coffee. In the estates sector, however, tank mixtures at half the normal rates, and copper-based fungicides at a reduced rate, are now widely used. The tank mixtures give effective control of CBD and result in good yields. They are less costly than the organic fungicides used by them, because of the reduced rates and the inclusion of the much cheaper copper fungicides. In

## FOREWORD

addition, the tank mixtures control coffee leaf rust and bacterial blight of coffee (two other serious disease of coffee) on the same spray programme.

With the present downward trend in coffee prices and increase in production costs, there is already a drastic decline in the profitability to the producers from coffee. The small-scale farmers lack capital and are not able to carry out in full, or at all, the recommended spray programmes. Coffee berry disease is therefore a constant threat to this major source of income.

The CRF realized as far back as 1967, that developing disease resistant varieties would be the long-term answer to the present problems. A breeding programme has been in progress at the CRF since 1971. Apart from CBD resistance, the programme has incorporated rust resistance improved yield, quality and compact growth in the new variety. The product of the CRF breeding programme is Ruiru 11, which was officially launched in 1985. Ruiru 11 is a hybrid arabica cultivar, which flowers within a year of planting and gives yield per tree, bean size and cup quality comparable to those of the established commercial varieties SL 28 and SL 34. The compact growth allows greater planting density thus increasing yield per unit area. To-date some 4,000 hectares of Ruiru 11 have been planted and it is expected to reduce the cost of production by about 30%. The mass adoption of Ruiru 11 has, in the past, been hampered by the low production of hybrid seed and seedlings; but the CRF is now producing more seedlings through vegetative propagation methods, and in the near future, through tissue culture.

In certain circumstances, the use of fungicides on coffee has been associated with increased coffee diseases, particularly CBD, in Kenya. Fungicides have also been reported to have 'tonic' effect on coffee, which increases growth and productivity above that expected from straight disease control. There is now some evidence that these effects might be mediated through changes in the microflora of the coffee surfaces. A recent study carried out at the CRF has established that specific types of microflora occur on coffee surfaces and these are affected quantitatively and qualitatively by changes in seasons, growth stages as well as fungicide regimes. The commonly occurring components of these microflora were shown to antagonize the CBD pathogen. The CRF hopes to pursue the possibility of using these natural surface microflora as a biocontrol method which can be integrated in the chemical control of CBD.

It is well known that research on CBD and other aspects of coffee is going on in respective African countries, perhaps more in some than others. However, since we have a common crop, coffee, with the same or similar problems, it would be to our mutual benefit to join our efforts and share our experiences. The network on CBD which has been constituted in Nairobi is intended to facilitate co-operative research in the effective and economical control of CBD in African countries. It is hoped that the network will solicit and co-ordinate funding for research programmes, bring together scientists to exchange experiences regarding the problem, as well as document and disseminate scientific literature within African. We therefore welcome the OAU's recommendation of constituting the network, thank the Inter-African Phytosanitary Council for sponsoring it, and Kenya for agreeing to host this important meeting.

**Wilson R Opilé**  
*Director of Research*  
*Coffee Research Foundation*

## **WELCOME ADDRESS BY HON. SIMEON NYACHAE, EGH, M.P. MINISTER FOR AGRICULTURE, LIVESTOCK DEVELOPMENT AND MARKETING**

Mr Chairman Distinguished Delegates, Ladies and Gentlemen,

On behalf of the Government of the Republic of Kenya, the Ministry of Agriculture, Livestock Development and Marketing, the Organizing Committee of the Constitutive Network meeting on the Coffee Anthracnose and myself, I would like to extend our warm welcome to Kenya to those that have come from other countries, and to you all the participants of this meeting. I hope your stay here will be pleasant and worthwhile. It gives me, my Ministry and the Coffee Research Foundation (CRF) great pleasure to host this very special meeting organized on coffee matters in Africa. We have been informed that the meeting was recommended by a joint Permanent Secretary set up by two resolutions (CM/RES 1173 (XL III) and CM/RES 1238 (E) of the OAU Council of Ministries in their 48th and 58th ordinary sessions.

Mr Chairman, it is common knowledge to us all that agriculture is the backbone of the economy of most African countries. Agriculture is also a way of life for the majority of our peoples, besides providing employment for a big percentage of our working population. For Kenya, this percent is 80%. For many of us, coffee is one of the principal agricultural export commodities. Until 1988/89, coffee was contributing on average 26% of the total domestic export, but its contribution has since dropped to 18% in 1990/91 as a result of the low prices experienced after the collapse of the price support mechanism of the International Coffee Agreement (ICA).

The importance of coffee to the national economy of Kenya convinced the Government, at the very early stage, of the need to have a centralized statutory organization which could implement the production and marketing policies in accordance with an Act of Parliament which was enacted in 1933. The Coffee Board of Kenya (CBK) is responsible for the running of the coffee industry broadly with respect to production, research, marketing, warehousing and advisory services to farmers and overseas promotion. The CBK has in turn delegated the research responsibilities to the CRF which is an official organization financed by the coffee farmers through the CBK.

Although coffee is an important commodity in our countries, we know that it has several problems: lack of proper know-how, especially at the small farmers' level who are the main producers; lack of adequate infrastructures like roads and warehouses; the rising costs of farm inputs like machinery and agricultural chemicals, drought or excess of rains now and then; pests and diseases; and in the recent past, low world market prices. These major constraints systematically require different and scientific approaches for their solutions. For many of these problems, strengthened research programmes and regional cooperation will be of vital importance. Coffee Berry Disease (CBD), the subject of this meeting, is a major problem in the African countries which produce arabica coffee; especially in Kenya, Tanzania and Ethiopia. As you may all know, CBD was first reported in Kenya in 1922 but has since slowly spread to Uganda, Zaire, Angola, Cameroon, Tanzania, Rwanda, Burundi, Ethiopia and more recently, to Malawi, Zambia and Zimbabwe. CBD has not been reported outside Africa and therefore, still remains solely an African problem. The losses due to the disease are enormous and despite successes in establishing and using fungicide control regimes, there are cases where crop losses of 20 - 30% are experienced during excessively wet weather conditions. Chemical control of the disease is expensive and imposes a continuing economic burden to the farmer. In Kenya, it is estimated that disease control contributes about 30% of the total cost of production of coffee.

Mr Chairman, with the present downward trend in coffee prices and increase in production costs, there is already a drastic decline in profitability to the producers from coffee. The majority of our smallscale farmers lack capital and are unable to carry out in full, if at all, the recommended spray programmes. An alternative long-term to the CBD problem is developing disease resistant varieties. Resistance to CBD is known to occur in some coffee varieties and has been used in Rwanda (Jackson Hybrid), Zaire (Bronze-tipped tyres), and Cameroon (Java variety). In Ethiopia, selection and multiplication of CBD resistant varieties from semi-natural coffee populations was made possible because it is the centre of genetic diversity of arabica coffee.

## WELCOME ADDRESS

In Kenya, the CRF opted to combine resistance to CBD and coffee rust with high yields and the good quality typical of Kenya coffee in a well-planned long-term breeding programme. The programme, which started in 1971, has produced a hybrid cultivar, Ruiru 11, which has been planted by Kenyan farmers since its official launching in 1985. This has been a milestone in the research on CBD in Kenya. Ruiru 11 is expected to cut down the cost of production by about 30%. The mass adoption of Ruiru 11 has, however, been hampered by the low production of hybrid seed and seedlings, and the high cost of establishment of the variety at close spacing. It is likely that screening of the wide variety of germplasm available in most of the African countries will lead to the eventual release of more CBD resistant cultivars. This is a long-term process requiring regional collaboration in germplasm exchange and selection for local adaptation.

It is well known that research on CBD and other aspects of coffee is going on in our respective countries, perhaps more in some than in others. However, since we have a common crop, coffee, with the same or similar problems, it would be to our mutual benefit to join our efforts at least at a regional level and share our experiences. We are all aware that there are few opportunities available to African scientists to meet and exchange technical findings, and information on our common problems even the circulation and distribution of scientific literature within Africa is so limited that very little is known of what is being done in neighbouring countries on similar problems.

In the area of CBD, however, there was a first regional workshop organized in Addis Ababa, Ethiopia, in July 1982. The workshop was organized by the Association for the Advancement of Agricultural Sciences in Africa (AAASA) in conjunction with the Ministry of Coffee and Tea Development and the Institute of Agricultural Research in Ethiopia and financed by the European Economic Commission (EEC). The workshop brought together delegates from many countries of the Eastern part of Africa and other eminent scientists from international organizations and institutions outside the continent. The proceedings for the workshop were published with the help of EEC and sent to individual participants. Amongst the resolutions and recommendations of this workshop was the need for exchange of information among African countries as well as the initiation of joint research programmes in neighbouring countries or those countries with similar ecological zones. The workshop was thought-of-as the beginning of an on-going regional CBD control development programme. Unfortunately, there has not been any follow-up meetings (over ten years now) and there is no way we can evaluate the progress made by individual countries regarding the technical resolutions and recommendations made. It is my hope that this Nairobi meeting is going to perform this onerous task.

More recently, the Inter-African Coffee Organization (IACO) has felt the need for improved research to increase the productivity and quality of coffee in Africa. The IACO has since formulated and constituted an African Coffee Research Network (ACRN also known as RECA in French), which is currently operative under the IACO Secretariat in Abidjan, Cote D'Ivoire. Kenya has been fortunate enough to have the Director of Research, CRF, elected as the first Chairman of the Network. The Network on CBD, which this meeting will constitute, shall, therefore, act as a specialist branch of the umbrella Network, ACRN. In this meeting, you are going to discuss in depth the current situation of CBD in our countries, the related research programmes, the results obtained, and their application at farm level. I hope at the end of your deliberations, you will come up with a stronger basis for the network and concrete plans for its future programmes.

Finally, may I take this opportunity to thank the General Secretariat, the Scientific, Technical and Research Commission and the Inter-African Phytosanitary Council of the OAU for having taken the initiative of organizing and sponsoring the meeting, and the CRF, in conjunction with my Ministry, for coordinating and hosting it.

Having said this, it is my pleasure and privilege to declare the meeting open and to wish you success in your deliberations.

Thank you!

# **STATEMENT BY DR. N. NKOUKA, AG. SCIENTIFIC SECRETARY OF THE INTER-AFRICAN PHYTOSANITARY COUNCIL OF THE ORGANIZATION OF AFRICAN UNITY (OAU)**

Your Excellency, the Minister for Agriculture, Livestock Development and Marketing, the Director of the Coffee Research Foundation, Honourable Delegates, Ladies and Gentlemen,

The subject matter which is uniting us here today in this beautiful country, Kenya, concerns a severe disease on Coffee (Anthracnose) in Africa. Dr Salim Ahmed Salim, Secretary General of the OAU attaches great importance on questions linked with food and agriculture, and in this context, I am highly honoured to represent him in this meeting. I also take this opportunity to transmit to you, the feeling of great importance which he is reserving to the outcome of this meeting of scientists, particularly, reinforced by various national experiences acquired in the field.

I do not want to hesitate in taking this opportunity, on behalf of His Excellency, the Secretary General of the OAU, to thank the Political Authorities of Kenya who did not spare a single minute to accept to host this important meeting here in Nairobi.

The high standard work already produced by the Coffee Research Foundation naturally weighed on this choice and the Director of Research, CRF, should kindly allow me to express here, on behalf of the OAU, my sincere thanks for their availability and the African warm welcome accorded to the participants.

Research scientists from various countries should transmit in a better way their experiences in order to avoid duplication, save more time and set up a more coherent African Scientific space. This is one of the goals of the Network on Coffee Anthracnose which we have just created in Nairobi. To create is one step but to ensure the care for breeding this baby which is just born will be thrilling and it is at that time that we shall need the experience of one another in order to make it come to our expectation.

Your Excellency, Ladies and Gentlemen,

I know that the task awaiting us is more important than this word of introduction, and while renewing the firm conviction of the OAU to give a helping hand to member countries to express their ambition for advancing research, I wish total success to the First Constitutive Network on Coffee Berry Disease (Anthracnose).

Long live the OAU, Long live the Inter-African Research!

I thank you.



## RESOLUTIONS AND RECOMMENDATIONS

The participants of the First Constitutive Network Meeting on Coffee Anthracnose wish to express their sincere appreciation and gratitude to the Organization of African Unity (OAU) in general and to the Inter-African Phytosanitary Council (IAPSC) in particular, for having conceived and sponsored this meeting. The idea of forming an Inter African research Network on the coffee anthracnose is indeed a profound positive step towards ensuring survival of the coffee industry, the back-bone of the economics of many OAU member states. The network shall pool together and utilise its own manpower resources, which have hitherto been working in isolation. To us participants, this meeting has been very educative and has given us the hope that the CBD problem can be overcome at a lower cost.

The participants express their utmost appreciation to the Government of the Republic of Kenya for hosting the meeting in Nairobi, and for the inspiring opening address by his Excellency, the Minister of Agriculture, Livestock Development and Marketing. They further acknowledge with gratitude the brotherly and warm welcome that was accorded to them during their stay in Kenya.

Participants would like to thank the Director of Coffee Research Foundation (CRF) for coordinating the meeting and also for the informative tour of the Coffee Research Station, Ruiru, which was conducted on the 28 May 1993. They learnt some valuable information which will help improve their own coffee programmes back home.

## CONSTITUTION OF THE NETWORK

It was agreed that the ability to directly sponsor national programmes on the coffee anthracnose is now limited because of the financial constraints individual countries are experiencing, and especially at a time when coffee as a commodity, is fetching very low returns. Funding organizations are also tending to sponsor joint or regional programmes as opposed to national programmes.

Since the Organization of African Unity (OAU) had offered to play a liaison role between member countries and financing Institutions, it was recommended that the formation of the Network on Coffee Anthracnose was essential. Participants were aware that the same OAU members through their membership to the Inter-African Coffee Organization (IACO) had recently formed the African Coffee Research Network (ACRN) which will deal with all research aspects on coffee. It was further explained that OAU through the Inter-African Phytosanitary Council, would play a liaison role between the proposed Network on Coffee Anthracnose and the ACRN to ensure that the two work in harmony.

## NAME OF NETWORK

The participants then resolved to form the network on the coffee anthracnose. The official name of "Coffee Anthracnose Research Network in Africa" (CARNA), was adopted.

## OBJECTIVES

The objectives of the Network are:-

1. To initiate and encourage formulation of joint research projects on the coffee anthracnose to member states
2. To document and disseminate information on coffee anthracnose within member states

## **RESOLUTIONS AND RECOMMENDATIONS**

3. To facilitate the exchange of research experiences between member countries through organizing scientific meetings, attachments, training, etc for researchers working on the Coffee Anthracnose.

### **MEMBERSHIP OF NETWORK**

It was resolved that membership be open to all African states (including Malagasy), where coffee is grown, irrespective of whether or not the coffee anthracnose is a problem.

### **HOME BASE**

It was resolved that Kenya, due to its more advanced research on the coffee anthracnose, become the home base of the Network. It was however, noted that the home base for CARNA could be moved to another member country in case of unforeseen problems.

### **COORDINATOR**

The participants of the 1st Constitutive Network meeting on Coffee Anthracnose in Africa, unanimously elected Dr (Mrs) Dinah M Masaba (Kenya), as Coordinator, and Dr Joseph Bakala (Cameroon) as Deputy Coordinator of CARNA.

### **FUNDING FOR NETWORK**

The Scientific Secretary (OAU) assured participants that the initial funding of CARNA will be undertaken by the Inter-Phytosanitary Council of OAU. It was hoped that later programmes could be maintained through external donor funding.

### **RECOMMENDATIONS**

After deliberating on the contents of the country reports and the discussions made during the different sessions of the meeting, the following recommendations were resolved.

### **CO-OPERATIVE RESEARCH PROGRAMMES**

The participants noted that individual countries were at different levels of research on the coffee anthracnose, and as such research priorities differ. It was, however, agreed that each country prepare sub-project proposals on areas of priority. The sub-projects were to be sent to the Coordinator who will draw out joint proposals representing the global needs on the coffee anthracnose. The joint proposals will be sent to the potential donors who will be identified by the Scientific Secretary (OAU). The proposal formats may vary depending on the donors. It was agreed that such formats as would be identified by the Scientific Secretary (OAU) should be sent to the Coordinator within the next three months after the meeting. It was also proposed that the Scientific Secretary (OAU) provide small grants to members to enable them prepare project proposals.

## EXCHANGE OF INFORMATION, MATERIALS AND PERSONNEL

It was noted that there were few opportunities, if any, available to African scientists to meet and exchange technical findings and information on common problems. Even the circulation and distribution of scientific literature within Africa is so limited that very little is known of what is being done in neighbouring countries on similar problems. Most countries were found to have a shortage of man-power and equipment for use on research on the coffee anthracnose, whilst there were hardly any exchanges in materials such as germplasm between countries. It was proposed that:

1. Each country start putting together all published information on coffee anthracnose; and individual lists be sent to the Coordinator for proper documentation and future dissemination within countries.
2. Each country gives an updated list of equipments and personnel working on the coffee anthracnose, giving details of academic qualifications, areas of specialization and deployment. The purpose of this exercise is to enable the network identify areas which require strengthening; organize training and research attachments/visits for scientists within member states; and identify expertise which can be used within the region.
3. Each country makes a list of available coffee germplasm to be sent to Coordinator for the purposes of future germplasm exchange programmes.

The above information in 1, 2 and 3 should be received by the CARNA Coordinator by the end of August 1993.

## INTERNATIONAL EXPOSURE

The participants, also noted that African Scientists will need exposure to International Scientific fora such as Conferences, Workshops, Symposia, etc which bear direct relationship to Plant Pathology in general, and to anthracnose diseases in particular, to enable them keep abreast with new technologies. It was recommended that CARNA undertakes to sponsor selected candidates to such meetings in future. In addition, CARNA will undertake to organize scientific meetings within member states, where scientists working on the Coffee Anthracnose will present and discuss their research findings.

## RESEARCH ACTIVITIES

From the discussions on the country situations of CBD, the following research activities were identified as needing priority attention:-

1. Most countries expressed the desire to screen germplasm for resistance to coffee anthracnose. The research would entail development and/or refinement of screening methodology. It was also noted that some countries, such as Kenya, were well advanced in breeding for resistance and the screening methods they use could be adopted and refined for use by the others. There was a feeling that the screening methods be standardized and the possibility of setting up joint screening facilities be explored.
2. Participants also identified a need for identification and documentation of pathogen variation (races) within member countries. The possibility of developing coffee varieties which would be used as differentials in race testing was suggested. It was also noted that there are other technologies such as the use of Restriction Fragment Length Polymorphism (RFLP) which could be explored for the same purpose.
3. Majority of countries were observed to be controlling the coffee anthracnose using fungicides. While it was agreed that individual countries carry out their own bio-efficacy evaluation, it

## **RESOLUTIONS AND RECOMMENDATIONS**

was felt that facilities for monitoring residual effects of pesticides (pesticide residue analysis) could be centralised to cut down on costs, especially because similar fungicides appear to be in use or under evaluation.

4. A report on the potential to control CBD using biocontrol was presented at the meeting. Participants recommended that research into new control measures such as biocontrol with surface microflora, or using plant extracts need to be actively supported and later incorporated into an integrated control approach to the coffee anthracnose.
5. It was observed that the distribution of CBD was not documented in some countries. Participants recommended that the spread of CBD in individual countries be monitored and documented. The spread of the coffee anthracnose to other African countries where it has not been reported, should be monitored and documented by CARNA.

## **PROCEEDINGS**

The Scientific Secretary (OAU) offered to publish the proceedings of the 1st Constitutive Network meeting on Coffee Anthracnose in Africa, in a special issue with assistance from the Coordinator.

## **CARNA MEETING**

It was resolved that in view of the urgency of the many activities which will need to be attended to within the starting period of CARNA, the next meeting of CARNA be held in one year's time i.e May 1994. The meeting will be held in Cameroon. The Scientific Secretary (OAU) is expected to invite potential donors to attend this meeting, where it is hoped that concrete project proposals will be presented and discussed.

Otherwise, members had agreed that CARNA should meet at least once every two years.

# COUNTRY REPORT: KENYA

D. M. MASABA, P. N. KING'ORI AND C. O. AGWANDA

*Coffee Research Foundation, P. O. Box 4, RUIRU, Kenya*

## INTRODUCTION

Coffee is a very important product in Kenyan agriculture, currently ranking third amongst the major sources of foreign exchange earnings. Up until 1988/89, the value of coffee represented 30% of the total domestic export, but the value has since dropped to 18% of the total domestic export in 1989/90 as a result of the low world market prices experienced after the suspension of the price support mechanism of the International Coffee Agreement (ICA) in 1989. Kenya coffee is produced by two Agricultural sectors, namely, the Plantation (Estates) and the Co-operative (Small-holders). The total area under coffee has increased over the years to the current figure of about 159,000 hectares. Total production has also generally increased over the years to a peak of about 129,000 tonnes clean coffee during 1987/88, but has since dropped to 87,000 metric tonnes in 1990/91 (Table 1).

Table 1: Coffee area production and export value in Kenya

Year	Area under mature coffee x 1000 ha			Production of clean coffee metric tonnes x 1000			Production per ha in tonnes clean coffee		Value of coffee		
	Small-holders	Estates	Total	Small-holders	Estates	Total	Small-holders	Estates	Mean price per tonne Kf	Total export value Kf x 10 <sup>6</sup>	% of total domestic export
1962/63	14.8	28.3	42.1	8.9	25.3	34.2	0.6	0.9	204	11.1	25
1963/64	25.7	28.3	54.0	15.4	28.5	43.9	0.6	1.1	350	15.4	33
1964/65	25.7	28.3	53.0	14.8	22.4	37.2	0.6	0.8	333	14.1	30
1965/66	36.6	32.0	68.6	25.6	25.7	51.3	0.7	0.8	327	118.8	32
1966/67	36.7	31.9	68.6	27.6	25.3	52.9	0.7	0.8	291	15.7	29
1967/68	49.9	31.2	81.1	20.6	13.3	33.9	0.4	0.4	320	12.8	28
1968/69	52.3	30.7	83.0	23.3	22.3	45.6	0.5	0.7	308	16.8	32
1969/70	54.1	29.9	84.0	26.3	26.5	52.8	0.5	0.9	373	22.3	37
1970/71	55.6	29.5	85.1	27.3	29.6	56.9	0.5	1.0	314	19.5	32
1971/72	55.7	28.7	84.4	29.4	31.9	61.3	0.5	1.1	375	24.8	27
1972/73	55.3	29.5	84.8	36.0	40.0	76.0	0.6	1.3	463	33.4	28
1973/74	55.6	29.1	84.7	40.6	32.7	73.3	0.7	1.1	510	38.4	24
1974/75	57.8	28.6	86.4	35.5	30.6	66.1	0.6	1.0	477	35.2	21
1975/76	57.3	28.6	86.4	36.5	38.1	74.6	0.6	1.3	1066	84.4	36
1976/77	56.6	27.8	84.4	37.7	49.7	87.4	0.8	1.8	2214	202.7	42
1977/78	65.0	30.6	95.6	47.6	33.7	81.3	0.7	1.1	1553	129.6	34
1978/79	70.0	30.0	100.0	46.7	27.6	74.3	0.7	0.9	1234	97.3	29
1979/80	80.0	30.0	110.0	52.0	39.7	91.7	0.7	1.2	1360	118.7	22
1980/81	84.7	32.9	117.6	64.0	34.7	98.7	0.7	1.0	1176	110.9	30
1981/82	75.5	33.6	109.1	52.5	34.4	86.9	0.5	1.0	1337	144.6	27
1982/83	101.0	33.6	134.6	52.5	33.0	85.5	0.5	1.0	1767	160.1	25
1983/84	114.2	35.7	150.0	74.7	54.3	129.0	0.7	1.5	2070	203.6	27
1984/85	116.3	35.7	152.0	67.7	28.9	96.6	0.5	0.8	2196	230.6	29
1985/86	117.7	38.6	156.3	68.4	45.5	113.9	0.6	1.2	2377	388.5	41
1986/87	116.0	38.5	154.5	67.9	36.4	104.3	0.6	1.0	2226	194.6	25.8
1987/88	116.4	36.7	153.0	84.4	44.5	129.0	0.7	1.2	2638	244.5	26.6
1988/89	116.1	39.6	155.7	78.3	38.6	116.9	0.7	1.0	2348	203.8	20.4
1989/90	117.5	38.1	155.5	69.5	34.4	103.9	0.6	0.9	1906	221.0	17.9
1990/91	121.5	38.0	159.3	51.4	35.2	86.6	0.4	0.9	2434	-	-

Figures from:  
 1. Coffee Board of Kenya Annual Reports  
 2. Central Bureau of Statistics, Kenya

The decline coincided with the unstable and depressed market situation resulting in low prices already alluded to above. This led to poor husbandry practices and generally low morale on the part of the farmers. The Smallholders contribute 64% while the Estates contribute 36% of the total production.

Kenya produces the mild Arabica coffee that is renown the world over for its fine quality. Coffee in Kenya is grown in areas between 1200 m and 2100 m above sea level. These areas lie on the broad gentle slopes of Mount Kenya and eastern Aberdare Range in the Central Province area, on the slopes of Mount Elgon, bordering Uganda, parts of the Great Rift Valley and some small holdings in the Taita Hills, a short distance from Tanzania (Fig 1).




Figure 1: Main coffee producing areas in Kenya

The production cost per tonne of clean coffee in Kenya is about Kf 2,700, which is considered high. About 30% of this total cost goes into the control of coffee diseases, the most important being coffee berry disease (CBD).

### COFFEE BERRY DISEASE

Coffee Berry Disease (CBD) attacks coffee berries in all stages of growth and occasionally even flowers and leaves. However, the most susceptible stages are during rapid berry expansion (6-16 weeks after flowering) and when the berry is ripening (Mulinge, 1970). On green berries, symptoms first appear as small dark, sunken patches which spread rapidly and may cover the whole berry (Fig 2a). Under wet conditions, pinkish mass of spores develop on the lesion surface. Usually, the fungus penetrates the interior and destroys the beans. Eventually, the whole berry dries out and takes the appearance of black mummified bodies locally known as 'mbuni' (Fig 2b). Infected berries may also be shed off as soon as lesions develop.




Figure 2a, b & c: Symptoms of CBD

A second type of lesion is normally formed under adverse weather conditions. This type of lesion is normally called 'scab' lesion (Fig 2c). Recently, it has been observed that some of the scabs are formed, especially on resistant coffee varieties even under weather conditions favourable to the disease and there is evidence that these scabs may be an indication of a defence reaction (Masaba and Van der Vossen, 1982).

## HISTORY AND SPREAD

The history and spread of CBD in Kenya has been documented in detail in Masaba *et al* (1984). Coffee berry disease was first reported in Kenya in 1922 (McDonald, 1926), on the slopes of Mt Elgon, but has since spread to all coffee growing areas. The disease is predominant in coffee growing areas above 1608 m in altitude, but occasionally becomes very severe in the lower altitudes. For example, during 1986/87, the disease was observed to be very high in the middle and low altitudes (Thika, Kilimambogo, Makuyu and Ruiru), prompting the CRF to carry out a survey to establish the possible reasons for the anomaly. It was observed that adoption of intensive irrigation practices had resulted into overlapping crops which may have led to the increase in CBD inoculum and the susceptible berry stages constantly being available on the trees, leading to enhanced CBD incidence (Mukunya *et al*, 1987).




Figure 3: The spread of Coffee Berry Disease across Africa



The spread of CBD to other African countries (Fig 3) has also been documented by Masaba and Waller (1992). The same authors observed that the spread of CBD is slow compared with the more rapid spread of the air-borne coffee rust. CBD spores are only released and dispersed by rain. They may, however, travel long distances on overtly diseased or latently infected plant material or passive vectors such as birds, man and machinery.

## THE PATHOGEN

The confused situation on the nomenclature of the *Colletotrichum* species on Arabica coffee was discussed in detail at the First Regional Workshop on CBD in Addis Ababa, Ethiopia, in 1982. The workshop recommended a critical review of the nomenclature of the CBD pathogen. The CBD pathogen had been given the name *Colletotrichum coffeanum* Noack (1901) which belonged to a saprophytic *Colletotrichum* species isolated on coffee from Brazil, where CBD does not occur. Subsequently, the International Mycological Institute (IMI) has carried out a detailed study aimed at distinguishing the CBD pathogen from other *Colletotrichum* species living on coffee and other crops.

The CBD pathogen habitually colonizes the maturing back of coffee shoots in close association with several other *Colletotrichum* species. However, only a small portion of the *Colletotrichum* spores produced from this source are pathogenic. Diseased berries are by far the major source of CBD inoculum (Gibbs, 1969). Fresh CBD *Colletotrichum* isolates can easily be distinguished from saprophytic *Colletotrichum* species by their distinct greenish grey cottony mycelia on malt extract agar in addition to their pathogenicity on green berries (McDonald, 1926; Rayner, 1952; Gibbs, 1969; Hindorf, 1970). However, after several subculturings or storage in culture, CBD isolates tend to revert to a white form indistinguishable from saprophytic *Colletotrichum* isolates (Waller, 1984). More recent investigations at the International Mycological Institute (IMI) have established a biochemical distinction between the coffee pathogen, *C. gloeosporioides* and *C. acutatum* which are the main saprophytic species on coffee (Waller, *et al*, 1993). CBD isolates cannot metabolize tartrate, or citrate as sole carbon sources, whereas the non-pathogenic *Colletotrichum* isolates from coffee, and also from other tropical woody hosts such as mangoes and citrus, can readily metabolize tartrate, citrate or both as sole carbon sources (Table 2).

Table 2: Features of *Colletotrichum kahawae* enabling it to be distinguished from *C. gloeosporioides* isolates from coffee

(1)	Colony characteristics (fresh single-conidial isolates on 2% MEA):
<i>Colletotrichum kahawae</i>	<i>Colletotrichum gloeosporioides</i>
Slow-growing (2-4 mm <sup>d</sup> 25°); profuse olivaceous to greenish dark grey mycelium; no acervular conidiomata produced; sporulation occurs from simple hyphae.	Faster growing (3-6 mm <sup>d</sup> 25°); white to pale grey mycelium; sporulation from acervuli or simple hyphae.
(2)	Metabolism
Cannot utilize citrate or tartrate as sole carbon sources.	Can utilize tartrate, citrate or both as sole carbon sources.
(3)	Pathogenicity
Pathogenic to young expanding green berries and developing seedling hypocotyls of <i>Coffea arabica</i> cv SL 28 and other susceptible cultivars, causing dark sunken anthracnose lesions.	Not pathogenic to young expanding coffee berries or seedling hypocotyls.

The results from the biochemical tests coupled with colony characteristics of fresh cultures on malt extract agar, and pathogenicity tests on expanding berries and seedling hypocotyls have formed a basis for recognizing the CBD pathogen as a distinct species named *Colletotrichum kahawae* Sp. Nov. The species name was derived from the Swahili and Arabic name for coffee, as CBD is still solely an African problem. The standard isolate used for description was from Kenya.

Recently, it has been reported that the Malawian isolates may vary in pathogenicity from the Kenyan isolates (Rodrigues *et al*, 1992) implying that there may exist physiological races of the pathogen.

## EPIDEMIOLOGY

Extensive research work was done on the CBD pathogen and disease in the late 1960s and 1970s and this has been reviewed by Firman and Waller (1977) and updated by Masaba *et al* (1984). It is particularly important to appreciate the central role of rain in spore production, dispersal, germination and infection by the pathogen, and also in regulating the flowering, and consequently cropping pattern of the coffee trees.

The coffee berry takes about nine months from flowering, which occurs shortly after the end of the dry season, to maturity (harvest). During the first four weeks, the berry does not increase in size, remaining at the 'pinhead' stage which is normally resistant to CBD. The expanding berry (4-16 weeks after flowering) is the most susceptible stage, while fully expanded green berries are resistant. The berries become susceptible again when they start to ripen (Mulinge, 1970).

Typically, in countries close to the equator, there tends to be two rainy seasons in a year. In Kenya, these are the long rains (March - June) and the short rains (October - November). The coffee growth pattern is thus characterized by two flowerings, one at the beginning of each rainy season, giving rise to two crops (the early short rains and late long rains crop) in each year (Fig 4).




Figure 4: Annual rainfall, coffee cropping patterns and spraying regime to control CBD in Kenya

## COUNTRY REPORT: KENYA

The crops thus overlap and hence provide a constant source of inoculum, which necessitates the application of fungicides throughout much of the year to protect the susceptible berry stages from infection. The cropping pattern is further complicated by fluctuations in the rainfall pattern and aberrant rainfall in the normally dry six months of the year. Irrigation during the dry season causes additional out-of-season flowerings which further exacerbate the overlapping of susceptible crop stages.

Some other factors influence the CBD epidemics. Among these are the frequent outbreaks of insect pests which attack the berry and increase its susceptibility to the disease. These include the berry borer (*Hypothenemus hampei* Ferrari) and the berry moth (*Prophantis smaragdina* Butler). Even where CBD is not present, the dead berries resulting from insect attack are often mistaken for symptoms of CBD by farmers.

### CHEMICAL CONTROL

The development of a successful fungicide control regime for CBD in Kenya has been exhaustively described in Masaba *et al* (1984). Spraying to protect the developing crop during the rainy seasons is necessary to achieve good control of CBD (Griffiths *et al*, 1971). Copper-based fungicides were used in Kenya until 1969 when Captafol and later Benomyl became available. The small-scale farmers in Kenya have continued using copper, mainly because of its relatively low cost, and also because copper gives effective control of other major coffee diseases namely coffee leaf rust caused by *Hemileia vastatrix* and in some areas, bacterial blight of coffee caused by *Pseudomonas syringae* pv *garcae*. In the estate sector, however, copper was almost wholly replaced by the more effective but expensive organic fungicides; captafol, chlorothalonil, dithianon, benomyl and carbendazim. After a few years of intensive use in Kenya, the pathogen developed resistance to carbendazim and benomyl and had to be withdrawn from the recommendations.

The fungicide-resistant strains are very stable and even without the selection pressure of the fungicide, they are still easily detected and have been found to interfere with control of CBD by contact fungicides (Masaba *et al*, 1990). The change from copper to organic fungicides also coincided with an upsurge of incidence of bacterial blight in high altitude areas within the Rift Valley (Okioga, 1978) as the bactericidal properties of copper are not shared by most organic fungicides and captafol exacerbated the disease (Kairu *et al*, 1984).

Table 3: Comparative chemical costs and yield benefits between control with straight fungicides and tank mixtures in Kenya

Chemical	Rate % formulated product	Clean coffee yield (kg/ha)	Yield benefit (kg/ha)	Cost of chemical per hectare (KSh)	Cost benefit ratio**
Captafol 80% WP	0.4	1169	735	6209	1:4.73
Captafol +	0.2)				
	+ )	1163	729	5567	1:5.24
Cupric hydroxide	0.5)				
Chlorothalonil	0.4	741	307	7920	1:1.55
Chlorothalonil +	0.2)				
	+ )	889	455	6209	1:2.93
Cupric hydroxide	0.5)				
Anilazine	0.4	852	418	5632	1:2.97
Anilazine +	0.2)				
	+ )	1011	577	5236	1:4.41
Cuprous oxide	0.5)				
Unsprayed	-	434	-	-	

\* Market prices for 1988;

\*\* Based on assumption that market price for coffee was K£2000/tonne but excluded labour costs.

Modified from Masaba *et al* (1990)

It was shown in Kenya that the use tank mixtures of organic fungicides at half the normal rates, and copper-based fungicide at a reduced rate of 5 kg/hectare, improves the cost/benefit ratio (Table 3). In addition, the tank mixtures are able to control coffee leaf rust on the same programme, and bacterial blight with a few additional copper sprays (Kairu, 1983). The tank mixtures are now widely used, especially on estates in Kenya. Tank mixtures constitute such a significant practice that some chemical manufacturers have formulated the components into a single mixture. However, these seem to be inferior to tank mixtures as they are only effective at higher doses (Masaba and Opilo, 1990).

Since its recommendation in 1969, captafol, and later its tank mixtures with copper, have consistently been the most effective fungicides giving comparatively higher yields than any other fungicide or fungicide combinations. Much of the captafol's yield benefit comes from its 'tonic' effect on coffee and persistence during the rainy season. However, captafol was recently banned from use due to its possible mutagenic effects. The removal of captafol greatly limits the options for fungicidal control of CBD. The current fungicides and fungicide combinations used for the control of CBD in Kenya are shown in Table 4.

Table 4: Tested and recommended fungicides for CBD control

Common Names	Proprietary Names (Trade Names)	Formulation	Rate (Kg/ha)
1. Chlorothalonil	Daconil 2787-W75	75% W P	4.4
	Clortocaffaro	75% W P	4.4
	Clortocaffaro 500 Flow	50% S C	4.4
	Jupital 720 Flow	72% S C	3.5
	Griffinil	50% W P	4.4
2. Dithianon	Delan	75% W P	3.3
3. Anilazine	Dyrene	75% W P	4.4
	Dyrene 480 S C	48% S C	6.0
4. Anilazine/Copper	Dyrene C 47	17/30 W P	9.0
5. Chlorothalonil/Copper	Dacobre 500	25/30 W P	7.7
6. Prochloraz-Mn	Octave	50% W P	2.2
7. 50% Copper formulations:			
(a) Cuprous Oxide (Red Copper)	(i) Copper Nordox	50% Cu W P	7.7
	(ii) Copper Sandoz MZ	.	.
	(iii) Copcel	.	.
(b) Cupric Chloride (Copper Oxychloride or Green Copper)	(i) Cobox	50% Cu W P	.
	(ii) Recop	.	.
	(iii) Microp-50	.	.
	(iv) Funguran	.	.
	(v) Cuprocal	.	.
	(vi) Copsap	.	.
	(vii) Cupravit	.	.
	(viii) Vitigram Conc	.	.
	(ix) Cuprado	.	.
	(x) Cuprocalfaro	.	.
	(xi) Vericuvre	.	.
	(xii) Kopox	.	11.0
	(xiii) Percopper	.	11.0
	(xiv) Blitox	.	7.7
	(xv) Cuprox	.	.
	(xvi) Oxycop	.	.
	(xvii) Cuprasol	.	.
(c) Cupric hydroxide (Blue Copper)	(i) Kocide 101	.	.
	(ii) Parasol	.	.
	(iii) Champion	.	.
8. Copper Sulphate Lime (Proprietary Pre-mix)	Frocida Bordeaux Mixture	25% WP	11.0

## BREEDING FOR RESISTANCE

The need to reduce the cost of production of coffee and particularly the cost of controlling coffee diseases, led to the initiation of a breeding programme in 1971. The programme aimed to combine resistance to CBD and coffee rust with high yields and the good quality typical of Kenya coffee (Van der Vossen and Walyaro, 1981). This was carried out in three stages:-

1. Selection within accessible variety collections followed by single crosses between disease resistant varieties and the best local cultivars (Table 5)
2. Multiple crosses to assemble in one plant the desired traits from several varieties
3. Backcrosses of selected plants from multiple crosses to the best local cultivars to improve quality (Van der Vossen and Walyaro, 1981)

Table 5: Yield and disease resistance for parent varieties and single crosses in Stage 1 of the Kenyan coffee breeding programmes

	Yield (first growth cycle) clean coffee tonnes/ha	Resistance in the field	
		CBD	Rust
<b>Parent varieties</b>			
SL28	10.9	S	S
Caturra	8.6	S	S
Padang	11.7	MR	S
Rume Sudan	2.5	R	MS
Hibrido de Timor	5.0	R	R
<b>Single crosses</b>			
SL28 x Caturra	11.4	Progeny screened for application of further selection criteria	
SL28 x Rume Sudan	8.3		
SL28 x Hibrido de Timor	12.8		
Caturra x Hibrido de Timor	10.3		
Caturra x Rume Sudan	9.7		
Padang x Rume Sudan	12.4		

The selection for resistance to CBD was greatly enhanced by the development of a reliable pre-selection test performed on six-week old seedlings (Van der Vossen *et al.*, 1976), which enabled large quantities of material to be screened under optimal laboratory conditions. The test was also used to identify the genes conditioning resistance in important progenitors. Rume Sudan (2 genes on the R- and K- loci) and Hibrido de Timor (1 gene on the T- loci), (Van der Vossen and Walyaro, 1980). The programme was then refined to ensure that all three genes were present in the new material. However, histological evidence suggests that the resistance mechanism is based largely on the formation of cork barriers, a type of resistance which is usually race non-specific (Masaba and Van der Vossen, 1982). Van der Graff (1981) showed that CBD-resistance in the Ethiopian selections was quantitative and probably polygenic, characteristics which also apply to resistant cultivars used in other countries. This information together with the lack of evidence for differential pathogenicity of CBD isolates and the continuing effectiveness of resistant varieties, provide some basis for expecting that the resistance may be durable.

In an exchange of germplasm in 1975 and 1977, the Kenya programme received some F<sub>3</sub> and F<sub>4</sub> selfings of Caturra x Hibrido de Timor (Catimor) from South America. The semi-dwarf Catimor hybrids have resistance to all races of coffee rust. The cross of Catimor with the selections of the main breeding programme resulted in the production of Ruiru 11 (Van der Vossen and Walyaro, 1981).

Ruiru 11 is a composite population of genetically heterogenous hybrids, though the plants show a high degree of phenotypic uniformity because of the dominance of the compact growth character derived from Catimor, and they all possess resistance to CBD and rust. Ruiru 11 flowers within a year of planting and gives yield per tree, bean size and cup quality comparable to those of the established commercial varieties SL 28, SL 34 and K7 (Owuor, 1988). The compact growth habit allows greater planting density thus increasing yield per unit area. To date, some 4337 hectares of close-spaced Ruiru 11 have been planted in Kenya. This represents approximately 6% of the national Arabica coffee tree population.

In order to develop varieties which possess all the three known resistant genes to CBD, a series of back-crosses and selfings are made annually. The resulting back-crosses or selfed seeds are reselected and planted out on preliminary trial fields for assessment of bean and liquor qualities as well as vigour. Concurrently, test-crosses are made on such genotypes in order to determine the number of CBD genes operating in them as well as their levels of homozygosity. The policy of the breeding programme at this stage is to retain only genotypes which have a minimum of two CBD resistant genes in homozygous condition.

Some 55 lines are currently being evaluated using the above criteria. Once the desired genotypes are identified, they will form the second generation of pollen supply parents (usually referred to as the male clones) in our hybrid seed production programme by 1995.

In order to completely restore the quality background of traditional Kenyan coffee varieties, a back-cross programme involving Catimor (currently used as the mother tree in the hybrid seed programme) and SL 28/K7 varieties was initiated in 1992. This programme is expected to introgress the desired compact growth of Catimor governed by the Ct-gene and the T-gene of resistance to CBD in this variety into SL 28 and K7 cultivars.

The search for more genes for resistance is also continuing in the CRS coffee gene banks. A number of introductions from Ethiopia are already being evaluated in a dialled cross in order to determine the number of genes controlling CBD resistance in this population. It is expected that stability of resistance to CBD will be greatly enhanced as more genes controlling resistance to this disease are identified and incorporated into future breeding lines.

## **BIOLOGICAL CONTROL**

In certain circumstances, use of chemicals on coffee has been associated with increased severity of coffee diseases in Kenya, particularly CBD (Furtado, 1969; Griffiths, 1972; Gibbs, 1972). Fungicides also have a 'tonic' effect on coffee, which increases growth and production above that expected from straight disease control. There is evidence that these effects might be mediated through changes in the microflora of coffee surfaces. The role of these surface microflora in the development of CBD in the field have been studied by Masaba (1991). Commonly occurring components of the microflora were tested in culture on detached berries and seedling hypocotyls for their effect on the pathogen. Several components were found to inhibit mycelial growth, spore germination and appressorial formation of the CBD pathogen (Masaba, 1991). It may now be possible to manipulate aspects of chemical control such as type of fungicide and application regime to favour the increase of populations of the antagonistic elements of the microflora.

## **CONCLUSION**

Despite earlier successes in establishing a fungicidal control regime for coffee berry disease, the disease continues to be a major constraint to coffee production in Kenya. Chemical control is expensive and imposes a continuing economic burden; furthermore its efficiency is frequently reduced by fluctuations in rainfall and cropping patterns.

The eventual replacement of Kenya coffee farms with the new disease resistant cultivar, Ruiru 11 is expected to have a considerable impact on the coffee industry, and the national economy as a whole. There will be a considerable reduction on production costs and improvement on cash incomes, especially for smallholders when they grow Ruiru 11. Ruiru 11 will also save the nation valuable foreign exchange spent on fungicides, fuel and spray machinery. Furthermore, the doubling in yield per unit area through close spacing of the compact Ruiru 11 variety will gradually release thousands of hectares of high potential land to food production. It is likely that screening the wider variety of germplasm available in most of the African countries will lead to the eventual release of more CBD resistant varieties and that is why regional collaboration is advocated.

The potential for biocontrol of CBD offers the development of integrated control packages in which an element of natural biocontrol is incorporated, though this area requires much more detailed research before it can become a reality.

It is important to stress that for efficient and economic control of CBD continuous research, coupled with collaboration in the form of exchange of information as well as materials, need to be strengthened. The CRF, therefore, fully supports the formation of a network on CBD in Africa.

## REFERENCES

- FIRMAN, I. D. AND WALLER, J. M. (1977). Coffee berry disease and other *Colletotrichum* diseases of coffee. *Phytopathological Paper No. 20*, Commonwealth Mycological Institute.
- FURTADO, I. (1969). Effects of copper fungicides on the occurrence of the pathogenic form of *Colletotrichum coffeanum*. *Transaction of British Mycological Society* 53:325-328.
- GIBBS, J. N. (1969). Inoculation sources for coffee berry disease. *Annals of Applied Biology* 64:515-522.
- GIBBS, J. N. (1972). Effects of fungicides on the population of *Colletotrichum* and other fungi on the bark of coffee. *Annals of Applied Biology* 70:35-47.
- GRIFFITHS, E. (1972). 'Negative' effects of fungicides in coffee. *Tropical Science* 14:79-89.
- GRIFFITHS, E., GIBBS, J. N. AND WALLER, J. M. (1971). Control of coffee berry disease. *Annals of Applied Biology* 67:45-74.
- HINDORF, H. (1970). *Colletotrichum* species isolated from *Coffea arabica* in Kenya. *Zeitschrift für Pflanzenkrankheiten Pflanzen Pathologie und Pflanzenschutz* 77:328-331.
- KAIRU, G. M. (1983). Integrated control bacterial blight of coffee and coffee berry disease. *Kenya Coffee* 48:103-109.
- KAIRU, G. M., NYANGENA, C. M. S., JAVED, Z. U. R. AND CROSSE, J. E. (1984). Iatrogenic effects of captafol on bacterial blight of coffee. *Plant Pathology* 33: 131-132.
- MASABA, D. M. (1991). The role of saprophytic surfaces mycoflora in the development of coffee berry disease (*Colletotrichum coffeanum*). *PhD Thesis*. University of Reading.
- MASABA, D. M. AND OPILO, V. L. (1990). The performance of formulated mixtures of copper and organic fungicides against coffee berry disease *Colletotrichum coffeanum* in Kenya. *Kenya Coffee* 55: 825-831.
- MASABA, D. M., KING'ORI, P. N. AND MUTHANGYA, P. M. (1990). Control of coffee berry disease (*Colletotrichum coffeanum*), using tank mixtures of copper and organic fungicides. *Kenya Coffee* 55: 949-956.

- MASABA, D. M., WALYARO, D. J. AND NJAGI, S. B. C. (1984). *Proc 1st Reg Workshop "Coffee berry disease", Addis Ababa 19-23 July 1992* Pg 125-130.
- MASABA, D. M. AND WALLER, J. M. (1992). Coffee berry disease. The current status. In *"Colletotrichum: Biology, pathology and control"*, Eds J A Bailey and M J Jeger Pg 237-249.
- MUKUNYA, D. M., ONDIEKI, J. J., OPILÉ, W. R. AND KAIRU, G. M. (1988). The current status of Coffee berry disease (CBD) occurrence and control in Kenya and factors associated with its increased severity. *Coffee berry disease Consultancy Report*, CRF.
- MULINGE, S. K. (1970). Development of Coffee Berry Disease in relation to the stages of growth of the berry. *Annals of Applied Biology* 65: 269-76.
- NOACK, F. (1901). Die krankheiten des kaffeebaumes in Brasilien. *Zeitschrift für Pflanzenkrankheiten* 11, 202.
- OKIOGA, D. M. (1978). The role of copper fungicides in the control of the berry disease. *Kenya Coffee* 43: 221-230
- OWUOR, J. B. O. (1988). An assessment of cup quality of the new disease resistant *Coffea arabica* cultivar Ruiru 11 in Kenya. *Kenya Coffee* 53 (619): 333-336.
- RAYNER, R. W. (1952). Coffee berry disease: A survey of investigations carried out up to 1950. *East African Agricultural Journal* 17: 130-158.
- RODRIGUES, C. J., VARZEA V. M. AND MEDEIROS, EMILIA F. (1992). Evidence for the existence of physiological races of *C. coffeanum* Noack *Sensu* Hindorf. *Kenya Coffee* 57:1417-1420.
- WALLER, J. M. (1984). Some mycological aspects of the coffee berry disease pathogen. *Proc 1st Reg Workshop "Coffee berry disease", Addis Ababa 19-22 July 1982* Pp 125-130.

## DISCUSSION

### E. Derso (Ethiopia).

- 1) What is the probable course of scab lesions?
- 2) Are there any races of *C. kahawae*
- 3) Is captafol still recommended for the control of CBD in Kenya?

### D. M. Masaba

- 1) Scab lesions on susceptible coffee varieties have been associated with unfavourable conditions for disease development. However, scab lesions have also been linked with host resistance (Masaba and Van der Vossen, 1982).
- 2) Captafol is no longer recommended for use against CBD in Kenya due to its alleged health hazard.

F. S. Ngulu (Tanzania). Can you elaborate on the biocontrol and how it is affected by use of fungicides?

D. M. Masaba. This was a wide project which established the possible role of saprophytic microflora to reduce CBD. Work concentrated on isolation and identification of the saprophytic microflora from coffee surfaces. The effect of fungicides used to control CBD on the microflora was only studied in



## COUNTRY REPORT: KENYA

field. We intend to study this interaction *in vitro* to establish specific effects of fungicides on individual microflora components.

Bakala (Cameroon). Why do you prefer use of fungicide mixtures to control CBD?

M. Masaba. Mixtures of copper and organic fungicides have been shown to be consistent in the control of CBD; cheaper at the rates used in the combinations; and can also control other coffee diseases such as *H. vastatrix* and *Pseudomonas garcae* on the same spray programme.

Sibomana (Rwanda). If Ruiru 11 is not stable, is it its propagation a problem?

O. Agwanda. Ruiru 11 is a hybrid with two selected parents. We use five mother parents and up to five male parents and that is why we say it is unstable. Otherwise its yield, quality, resistance to CBD and leaf rust, and its compact growth characteristics are very stable.

## COUNTRY REPORT: TANZANIA

### RESEARCH SYSTEM ON COFFEE BERRY DISEASE AND ITS CONTROL IN TANZANIA

F. S. NGULU

*Lyamungu Research Institute, Moshi*

#### ABSTRACT

Coffee is the major foreign exchange earner in Tanzania; Arabica coffee accounts for over two thirds of the total coffee output. Coffee Berry Disease (*Colletotrichum kahawae*, Waller & Bridge) spread to all major coffee producing areas within 14 years since it was first reported in the country in 1963. It is of economic importance on farms situated 1300 m.a.s.l. CBD control research programme consists of inoculating and assessing breeders' material for resistance as well as testing the efficacy of fungicides against the disease under field conditions. Regional/International cooperation would consolidate research efforts through germplasm exchange, standardizing research techniques, monitoring the effects of long term fungicide use and sharing research findings through exchange of reports, meetings and field visits.

#### INTRODUCTION


Coffee is currently the leading foreign currency earner in Tanzania. Annual coffee production is approximately 55000 MT of which 2/3 is Arabica and the remainder is Robusta coffee. The major Arabica coffee producing areas are Arusha and Kilimanjaro in the North-East and Ruvuma and Mbeya regions in the Southern highlands. Robusta coffee is grown mainly in Kagera region on the Western shores of Lake Victoria.

Production of coffee is mainly by small holders, owning between 0.5 - 2.0 hectares and they account for 95% of the total output. On small holdings coffee trees are mostly capped; either as single stem, multiple stem, interplanted with bananas & occasionally with shade trees such as *Maranguensis* and *Grevillea robusta*.

Current commercial varieties of Arabica coffee were derived from Bourbon and Kent selections. They are high yielding varieties with good liquoring quality but are susceptible to the two major diseases; i.e. Coffee Berry Disease (*Colletotrichum kahawae* Waller & Bridge) and Coffee Leaf Miner (*Hemileia vastatrix* Berk & Br.).

Coffee Berry Disease (CBD) was recorded for the first time in Tanzania in Tarime district, (Kibani, 1963 and Bujulu, 1974), an isolated coffee growing area in North Western area of the country. The first CBD incidences in the main arabica coffee producing areas in Kilimanjaro/Arusha and Southern highlands were reported in 1966 and 1977 respectively (Bujulu, 1974 and Kibani, 1977). In other words, geographical isolation notwithstanding CBD spread to all major coffee growing areas within 14 years after it was first reported.

The disease is of economic importance in coffee farms situated above 1300 metres a.s.l. where losses on individual farms range from 30 - 60% depending on weather conditions particularly rain.



- Coffee growing areas
- International boundaries
- + CBD infected areas

Figure 1: Coffee producing areas in Tanzania and CBD incidence

## COFFEE RESEARCH PROGRAMME

The programme consists of five disciplines, viz. Plant Breeding, Plant Nutrition, Agronomy, Entomology and Pathology. Like the other crop research programmes coffee research is under the Ministry of Agriculture, Livestock and Cooperatives. It is headed by a National coffee research coordinator who is appointed by the Ministry. Under the Coordinator is a team of researchers who are required to draw up research project proposals in their respective fields of specialization. Research proposals are submitted to a technical committee which meets once per year. Research findings on approved projects have to be presented to the same committee for ratification. Subsequently, approved research findings are communicated to the Extension services and agricultural research institutes in the form of Annual reports; seminars and field days. The programme has a specific training wing.

Projects on CBD control fall under two categories viz. screening breeder's material for resistance and testing the efficacy of candidate fungicides against the disease under field conditions.

Initially, screening of breeder's material was limited to the hypocotyl preselection test as described by Van der Vossen *et al* (1976). However, lately, this exercise has been expanded to include assessments on individual bearing trees in the field. In the latter case, a branch with suitable berry count at the most susceptible stage (soft-green) is enclosed in polythene sleeve and sprayed with an inoculum in a manner similar to the hypocotyl test. The level of resistance of the individual tree relative to that of a susceptible control is determined by expressing the infected berry count as a percentage of the total count. To assess the level of resistance of hypocotyls, an infection scale (see appendix 1) instead of 1 - 12 as described by Vossen *et al* (1976) is used.

Testing the efficacy of candidate fungicides against CBD is carried out under field conditions. A set of candidate products is usually tested for 3 crop seasons in 2 - 3 different sites. Manual Knapsprayers are used to apply the treatments. Gross plot size commonly used is 25 trees (5 x 5) with a net plot of 3 x 3 trees. On each tree in the net plot, single branches at top, medium and low elevations are marked prior to CBD onset. These branches are subsequently assessed monthly for natural infection. The number of infected berries is expressed as a percentage of total berry count.

It is worthy noting at this stage that all fungicides tested to-date for CBD control are essentially "protectants" and recommendations issued to coffee growers require them to apply the fungicides repeatedly at specified intervals throughout the period when a large proportion of their coffee bushes are vulnerable to infection. The major snag with the above recommended package is that in years of low rainfall where 150 - 200 mm ten-day rainfall total is annually recorded, it becomes difficult and expensive to sustain fungicide deposit on coffee bushes and the efficiency of disease control is disrupted.

## LIMITATIONS AND NEEDS

Inadequate funding particularly foreign currency to replenish consumable laboratory items and spare parts for laboratory equipment and procurement of relevant journals.

An isolated working environment. There exists no formal forum where coffee scientists can share their experience with their counterparts outside the country.

Unsatisfactory level of manpower strength and continuity. Coffee being a perennial crop requires assured continuity of the services of qualified scientists so that cost-effective packages for control of CBD adoption can be realised in the shortest possible time period.

## REGIONAL/INTERNATIONAL COOPERATION

Regional/International cooperation would be highly desirable particularly with regard to germ-plasm exchange, standardizing research techniques, exchanging research information and field visits with the aim to minimize chances of work duplication on common research problems.

## LITERATURE

ANONYMOUS (1963). *Coffee Census up to December 1963* North Mara.

NGULU, J. (1974). *Historical background of Coffee Berry Disease in Tanzania*. Lyamungu progress report.

NGULU, J. (1980). *Miscellaneous report on Coffee Berry Disease*, Lyamungu.

VAN DER VOSSEN, H. M. COOK, R. T. A. and MURAKARU, G. M. W. (1976). Breeding for resistance to Coffee Berry Disease caused by *Colletotrichum coffeanum* in *Coffea arabica*. *Euphytica* 25: 733 - 745.

## DISCUSSION

M. Masaba (Cameroon). Could you give more information on hypocotyl inoculation?

J. Ngulu. The method was developed in Kenya (Van der Vossen *et al*, 1976) but we have just modified the scale of recording. We use five instead of the original 12 mean grades.

M. Masaba. At what age do you inoculate your berries?

J. Ngulu. At 14 weeks after flowering, which is still in the susceptible expanding stage.

M. Masaba (Kenya). How do you control the weather factor during field inoculations when you find the weather at Lyamungu not being favourable for CBD development?

J. Ngulu. We usually inoculate in April, May when the weather is ideal.

M. Masaba. How about using the hypocotyl technique for your resistance screening?

J. Ngulu. We noted susceptibility in some lines which were passed as resistant by hypocotyl technique.






M. Mwang'ombe (Kenya). You have reduced the scale to 1 - 5, how do you rate?

J. Ngulu. 4 - 5, very susceptible; 1 - 3, the material goes straight into the field.

# COUNTRY REPORT: TANZANIA













## APPENDIX I:

### HYPOCOTYL INFECTION: ASSESSMENT SCALE 1 - 5

Seedling appearance	Scale	Symptom description
	1	No visible disease symptoms.
	2	Few to several brownish lesions. Tiny to enlarging but non-expanding.
	3	Small sunken dark lesions. Starting to expand laterally and vertically.
	4	Large coalescing or individual sunken lesions. Seedling not shrivelled.
	5	Large and sunken. Seedling completely girdled, shrivelled and dead.

ENDIX II:

POCOTYL INFECTION: ASSESSMENT SCALE 1 - 12

<p>1</p>  <p>No visible symptoms</p> <p>R</p>	<p>2</p>  <p>A few scab lesions</p> <p>R</p>	<p>3</p>  <p>Small scab or tiny brown lesions</p> <p>R</p>	<p>4</p>  <p>Scab or brown lesions</p> <p>R</p>
<p>5</p>  <p>Scab &amp; brown lesions &amp; a few small black lesions</p> <p>MR</p>	<p>6</p>  <p>Brown and narrow black lesions</p> <p>MR</p>	<p>7</p>  <p>Narrow black lesions some &gt; 1 cm long</p> <p>MR</p>	<p>8</p>  <p>Black lesions becoming wider &amp; starting to coalesce</p> <p>MR</p>
<p>9</p>  <p>Large coalescing black lesions but not yet complete girdling</p> <p>S</p>	<p>10</p>  <p>Large coalescing black lesions complete girdling of stem</p> <p>S</p>	<p>11</p>  <p>Most of the stem affected &gt; 1/3 stem shrivelled seedling dead</p> <p>S</p>	<p>12</p>  <p>Whole stem affected and shrivelled seedling dead</p> <p>S</p>

## COUNTRY REPORT: ETHIOPIA

### REVIEW OF THE CURRENT STATUS OF COFFEE BERRY DISEASE (CBD) RESEARCH AND DEVELOPMENT IN ETHIOPIA

ESHETU DERSO

*Institute of Agricultural Research Jima Research Centre, P. O. Box 2003, ADDIS ABABA, Ethiopia*

In Ethiopia, *Coffea arabica* L. grows in almost all regions with conditions ranging from the semi-savannah climate of the Gambella plain (1500 m altitude a.s.l.) to the continuously wet zone of the south-west and in gardens and back yards (1500 - 2600 m altitude a.s.l.). The soil ranges from sandy loam to heavy clay while the general soil types which are acidic (pH 4.2 - 6.8) are reddish-brown lateritic loams or clay loams of volcanic origin. The annual rainfall varies from 1000 to 2400 mm (Sylvain, 1958; Fernie, 1966).

From a genetic point of view, arabica coffee is the most important crop of the nation. Almost the entire genetic diversity is found mainly in the south-western rain forest area where almost undisturbed patches of forest with coffee as undergrowth can still be found.

Coffee grows as shrub or small tree and as a semi-wild crop in the moist Montane forest. As a commercial crop it is cultivated under ranges of shade conditions and where grown under irrigation in dry areas, it is grown without shade. As a garden plant, it is often mixed with ensate fruit trees and herbs in the backyard (Tewolde and Gebre, 1990).

The area occupied by coffee varies from 500,000 to 550,000 ha and the bulk of the production comes from poorly managed forest coffee. Coffee is the major cash crop in Ethiopia. It contributes 5% of GDP, 12% of the Agricultural sector output, 70% of the foreign exchange earnings, 10% of the government revenue and employs 25% of the workforce of the country (Mesfin, 1989).

A major research effort was begun by the National Coffee Research Centre (NCRC) which is part of the Institute of Agricultural Research (IAR). The Centre was established in 1966 to formulate the National Coffee Research Policy and conduct and coordinate research in the Country.

The NCRC conducts its coffee research activities at Melko, the main research centre and at sub-centres namely Agaro, Gera, Metu, Tepi, Anfilo, Wonago and Mechara which are located in different coffee growing regions of the country.

The Institute of Agriculture Research has set a national coffee commodity team composed of members from the national centre, cooperating members from IAR and Non-IAR research centres and invited scientists or subject matter specialists from user organizations. The team reviews the relevance and scientific quality of new proposals and monitors the on-going programmes. The team also has the mandate to accept or reject new proposals on scientific grounds.

Coffee research priorities are always determined by scientists in close cooperation between the Ministry of State Farms Coffee and Tea (MSFCT) and the Coffee research team and thereafter approved by the Institute of Agricultural Research (IAR). Experiments are also conducted in the sub-centres under the supervision of the Jima National Coffee Research Centre. To adapt or develop and transfer improved coffee production technologies, the National Coffee Research Centre plays a leading role. Coffee research goals are defined clearly by NCRC in light of the mandate of the MSFCT, which is the only responsible organ for the promotion of our number one foreign-currency earner crop (Mesfin, 1989).



Coffee Pathology Section consists of 15 personnel among which two are research officers, one assistant research officer, one Junior research officer, two laboratory technicians, four technical assistants and four support staff.

Laboratory work with disease causing organisms need meticulous handling of the pathogens and the exemplars and this is, of course, subject to the facilities available in the laboratory and the quality of technical knowledge available. Obviously, the dearth for the above mentioned infrastructure becomes a limiting factor to the dissemination of new technologies to users. Currently, the section is equipped with seed germinators, moist chambers, oven, centrifuges, incubators, desiccators and refrigerators (with alternating temperature).

At the moment, even though the documentation system of the Institute of Agricultural Research is based on electronic data processing technology, the section lacks computer data base system and also lacks adequate well trained and experienced personnel. The section is still confined to the traditional system of filing and computing where each data is coded in a file and placed in a filing cabinet.

Modern computer data handling facilities are essential for the Pathology Section for efficient and effective integration of information and data collection from all relevant sources and disciplines.

Some of the coffee research activities are resource demanding and are too costly to be handled. External support will therefore be needed, especially for new areas or those that represent expansion of on-going activities based on pertinent national policy. Such support should be sought through collaborative links with regional and international institutions working in the field. In this regard, an agreement has been reached between IAR and the European Economic Commission (EEC) to employ an internationally competent high calibre Plant Pathologist and two scientists in other fields, for three years, who would assist and guide coffee researchers with their research activities.

As far as training of the Coffee Pathology section staff is concerned our conviction is that every one in the field assistant to research officers need training and that the training of every one is of equal importance. Training is a success and very vital for all involved in research activities and must be a continuous process, it can never be static. To our dismay, training in the NCRC remained a bread and butter issue which has got no remedy. Therefore, the assistance of international institutions in the field is of paramount importance.

Plant material distribution and exchange is performed by the crop protection and regulatory department of the Ministry of Agriculture (MOA) which issues plant importation permits to relevant applicants. The department checks for diseases and pests on the incoming material. Material which is dispatched abroad is also certified. The IAR quarantine Officer, on behalf of the IAR staff is responsible for issuing the permit for material importation and exportation.

## **CURRENT STATUS OF CBD IN THE COUNTRY**

Coffee Berry Disease (CBD) is an anthracnose of the green and ripening berries and causes very heavy crop losses in arabica coffee if not controlled. Yield losses caused by this disease remain among the major constraints to increased coffee production in many parts of Ethiopia.

When CBD was first identified in Ethiopia in 1971, it rapidly spread throughout the country and caused a great concern in coffee production. Currently according to five years mean result (1987 - 1991) at Gera, the loss due to this disease exceeds 40%. However, the overall national loss in the country is estimated to be between 25 - 30% (Eshetu and Girma, 1992).

Since the first appearance of this disease in Ethiopia, measures have been taken to keep the disease intensity below the level at which it may become economically important. Accordingly screening of coffee genotypes against the disease was considered as a short term control measure, while the development of CBD resistant coffee types was taken as a long term control measure.

## CHEMICAL CONTROL AGAINST CBD

In any agricultural system, chemical means of control should be considered as a last resort and an alternative to the other principles of plant disease control. However, if this control measure is planned, it must be borne in mind that it should be carried out sensibly, taking into account economic and ecological considerations.

Coffee cultivars grown by the State Farms in Ethiopia are all CBD resistant and also there are hundreds of hectares of peasant holdings planted with the resistant varieties. These farms, however, constitute a very small proportion of the total area cultivated under coffee. On the contrary, the total production in the country comes from the peasant sector which consists of mainly garden and semi-forest coffee, with variable degree of susceptibility. Hence, it goes without saying that chemical protection against CBD in our time is indispensable.

Besides, chemical control is well suited to coffee because the value of coffee allows the cost to be economical. Other methods of control which rely on changing crop varieties, crop rotation and cultivation have not been used to any great extent. However, the effect of certain pesticide products on the organoleptic qualities of coffee need another consideration.

As a short term means of control, fungicide screening for the control of CBD in Ethiopia started as early as 1972. Since then attempts have been made to select different fungicides. Currently, it is considered important to continuously screen fungicides against CBD in order to provide growers with an ample choice of fungicides that are effective, safe and cheap (Eshetu, 1990).

Among the various fungicides which, from time to time, have been subjected to experimentation against CBD at Gera (Table 1), some have shown good results both in controlling the disease and in yield increase. Therefore, their use in the country against CBD has been encouraged.

Table 1: Tested and recommended fungicides against CBD in Ethiopia

Common Names	Trade Names	Formulations	Rate Kg/ha L/ha
Chlorothalonil	Daconil 2787	75% W P	4.4
Prochloraz	Octave	50% W P	2.0
Chlorothalonil	Griffiniil	75% W P	4.4
Cuprous Oxide	Nordox	50% W P	7.7
Prochloraz Copper Oxychloride (Proprietary Pre-mix)	Octave Super	50% W P	6.5
Chlorothalonil + Cuprous Oxide (Tank-mix)	Daconil + Nordox	75% W P + 50% W P	7.0

Apart from these, there were some fungicides which have earlier been used in the country against CBD, but they have been withdrawn from use due to the reported carcinogenic effects, obsolescence and phytotoxicity.

## DEVELOPMENT OF CBD RESISTANT COFFEE TYPES

diversity of *Coffea arabica* L. can be considered as a corner stone for selection of resistant coffee and this has given the country the chance to have a competitive advantage to develop genotypes resistant to CBD and which can best adapt to particular regions.

At the out-break of CBD, significant achievement has been made in selecting resistant cultivars to CBD and by the end of the fifth year of research, over 15 selections were identified possessing high levels of resistance to CBD. From these selections, millions of seeds were released for planting and the level of resistance of their progenies under plantations were found to be as stable as their parents (Eshetu and Bayeta, 1982).

As part of the major selection programme for CBD resistant coffee types in 1973, a number of selection and collection programmes have been conducted in the eastern, southern and south-western growing regions of the country. These new selections are planted in progeny plots at different localities under conditions of heavy CBD infection so that they can be evaluated under different environmental conditions which prevail in the areas of potential use of the selections and collections for their resistance to CBD, agronomic performances and other essential characters. Moreover, these selections were also being investigated in the laboratory at the NCRC for their level of resistance to CBD using inoculation of detached berries. The main purpose of widening the scope of selection and collection programmes is to avoid adaptation problems and the threat of losing potentially useful coffee types which might be realized if the heterogenous forest coffee is replaced by few resistant coffee types selected from specific localities or regions.

Not least, there are some points of emphasis regarding CBD research in Ethiopia and these are:

Control of CBD should be directed towards an Integrated Pest Management (IPM) strategy. A strategy which entails the simultaneous or sequential use of several methods of control. Biological control as a component of IPM is of particular importance. To fulfil this objective regional and international cooperation should be sought to consolidate CBD programmes financially and with skilled and adequate manpower.

The significance of Ethiopia as a source of important diversity in Arabica coffee can be illustrated by the germplasm flow in and out of the country since historical times. But at the moment, international exchange of seeds and propagation materials to and from Ethiopia needs special consideration since it may pose a serious problem of pest introduction and dissemination risk as the quarantine service in the country lacks essential elements that would enable it to apply appropriate measures.

Interactions between and amongst scientists are crucial to the generation of new ideas and to the best possible exploitation of existing knowledge. Every effort should be made to foster mutual understanding and trust between scientists in different disciplines in Africa.

## REFERENCES

ESHETU, D. and DERSO. *Evaluation of fungicides for the control of CBD in Ethiopia*. p. 85 - 87. In: Seid Ahmed and Yaynu Hiskias (eds) 1991. Proceedings of the 15th Annual Meeting of the Ethiopian Phytopathological Committee. 13 - 14 March, 1990, Addis Abeba, Ethiopia.

ESHETU, D. and GIRMA, A. *Determining level of CBD resistance on progenies of 1982 selections by the detached berry test*. p. 50. In: Eshetu B. et al (eds) 1992. Proceedings of the joint conference Ethiopian Phytopathological Committee and Committee of Ethiopian Entomologists. 5 - 6 March, 1992, Addis Abeba, Ethiopia.

FERNIE, L. M. (1966). Some impression of coffee in Ethiopia. *Tanganyika Coffee News of July 11*. pp. d.

MESFIN, A. and BAYETA, B. (1982). *Resistance of the F1 to CBD in six parent diallel cross coffee*. In: Proceedings of the first regional workshop on Coffee Berry Disease. 19 - 24 July, Addis Abeba, Ethiopia.

MESFIN, AMHA (1989). *Managing coffee research in Ethiopia*. Paper presented to the IAR research management workshop 23 - 28 January, 1989. Debre Zeit, Ethiopia.

TEWOLDE BERHAN, G. E. *The importance of Ethiopian forests in the conservation of arabica gene pools*. Paper presented to the 12th plenary meeting 4 - 10 September, 1988. Hamburg, Germany.

SYLVAIN, P. C. (1958). Ethiopian coffee: its significance to world coffee problems. *Econ. Bot.* - 139.

## DISCUSSION

D. M. Masaba (Kenya). You talked of coffee grown in gardens and forests, can you explain this?

E. Derso. Small scale farmers usually have coffee which is still under forest (semi-forest) or in gardens. Under this situation, farmers are not allowed to interfere with the habitat, but they are allowed to use fungicides to control CBD.

C. O. Agwanda (Kenya). How come small scale farmers grow susceptible coffee and government estates grow resistant lines.

E. Derso. Resistant lines are selected from the semi-forest coffee and passed on to government estates to grow.

C. O. Agwanda. To protect the forest coffee, the small scale farmers should be supported to help them conserve the germplasm, especially in the light of the low coffee prices.

E. Derso. This is already being done.

P. Zvoutete (Zimbabwe). Is there any work that has been carried out to identify races of the pathogen?

E. Derso. None.

N. Nkouka (OAU). How are the plant quarantine regulations in Ethiopia?

E. Derso. At the moment not clear regarding coffee.

A. W. Mwangombe (Kenya). What technique do you use to screen coffee lines for resistance to CBD?

E. Derso. We use detached berry test in the laboratory and attached berry test in the field. Hypocotyl screening test was not found to be consistent in Ethiopia.

D. M. Masaba (Kenya). This is contrary to the results and experiences in Kenya as we have consistently found the hypocotyl technique to be more correlated to attached berry test, and detached berry test to be very inconsistent in infection results.

# COUNTRY REPORT: ZIMBABWE

## COFFEE BERRY DISEASE IN ZIMBABWE

P. ZVOUTETE

*Plant Protection Research Institute, P. O. Box 8100, CAUSEWAY, Harare*

### CBD OUTLOOK

Coffee Berry Disease was first reported in Zimbabwe in 1985 and has now been confirmed on 10 of the thirty coffee farms in Mashonaland, West Province in Northern Zimbabwe. The main coffee growing areas in the Eastern Highlands which account for 90% of the country's coffee production are at the moment free from the disease. The main coffee variety grown, cv SL 28, is susceptible to CBD. The other varieties cvs Caturra and Catimor, are moderately resistant. The disease was recorded in 1991/92 not being serious during the last two growing seasons (1991/92) because of the drought the country experienced. Yield losses vary from year to year and from plantation to plantation depending on the age of the crop, management, variety grown and prevailing climatic conditions. Yield losses due to CBD vary from less than 10% to more than 60% in some plantations.

The strain of the causal fungus, *Colletotrichum kahawae* in Zimbabwe appears to be identical to the one elsewhere in Africa according to tests carried out by IMI.

There are a few fungicides commonly used by farmers to control CBD in Zimbabwe and these are anilazine (Dyrene), Captafol, Copper Oxychloride and Chlorothalonil. Within the country, quarantine measures are in place to regulate coffee movement from the CBD infested areas.

### NATIONAL COFFEE RESEARCH PROGRAMMES

Coffee research is carried out at Chipinge Research Station in Chipinge and on some private coffee farms. Plant protection aspects are researched on by Plant Protection Research Institute. There are also some agro-chemical companies who screen pesticides for control of coffee plant diseases. Chipinge Research Institute carries out physiological and agronomic research. The Coffee Growers Association (CGA) also supports research on coffee.

Current general coffee research programmes include the following:

#### A. Agronomy/Physiology Programmes

1. Irrigation trials
2. Planting depth trials
3. Variety trials
4. Row orientation trial
5. Plant growth regulators (PP.333)

## **Plant Protection Programmes**

1. Chemical control of Antestia bugs
2. Leafminer insecticide trial
3. Leaf rust fungicide screening trials
4. CBD fungicide screening trial
5. Study of the effect of canopy on CBD development and dispersal

## **PERSONNEL WORKING ON CBD**

There is a Senior Research Officer working on CBD but he is currently on study leave. Other biologists at PPRI assist in CBD research. There are two Research Technicians working on CBD. A plant pathologist was recently engaged and will be based at Chipinge Research Station. She is expected to take up all responsibility for coffee disease including CBD. There are crop protectionists and agro-chemical companies screening fungicides against the disease.

## **INFRASTRUCTURE**

Facilities at PPRI used for pathological research on other crops are also available for CBD work. Most of the work is carried out on private farms in the CBD quarantine area and the land is usually worked with assistance of CGA. The work is financed principally through a government vote to PPRI for plant disease research. CGA periodically supports some projects in coffee.

## **RESEARCH STATIONS**

The bulk of coffee research is done at Chipinge Research Institute in Chipinge in Eastern Zimbabwe. However, there is no CBD research at CRI because the disease has not been reported in this region. Field work on CBD is carried out mainly at Magunje Sub-station while laboratory work is at PPRI in Harare.

## **DOCUMENTATION FACILITIES**

All personnel have easy access to university and other libraries but these institutions do not have extensive literature on coffee. CGA also provides information bulletins to PPRI on request. PPRI has links with IMI in UK; CIFC in Portugal and Coffee Research Station in Kenya.

## **SCIENTIFIC EXCHANGE**

There are negligible regional scientific exchanges, plant material exchanges and any other form of cooperation with PPRI. There are some plant material exchange between CRI and coffee research stations in the region.

## **TRANSFER OF SCIENTIFIC RESULTS**

Scientific data generated by PPRI and indeed by the whole Department of Research and Services (DR&SS) is transferred to the user communities principally through an agricultural extension department, the Department of Agricultural and Technical Extension (AGRITEX). There are committees comprising the two departments which meet periodically to review agricultural extension programmes. PPRI researchers conduct training courses for AGRITEX staff to update them on pests and diseases. The different institutes under DR&SS produce annual reports which are distributed to AGRITEX's regional offices and any other interested bodies. Plant protection information is also disseminated through PPRI sub-stations which are located in some of the coffee growing areas. There are also periodic meetings with commodity based associations such as the Coffee Growers Association.

Although the structures exist for dissemination of data, in practice the coordination is weak and the impact is therefore less than can be achieved.

## **CONSTRAINTS AND NEEDS IN CBD RESEARCH**

Agricultural research in Zimbabwe is bedeviled at present with many limitations. The major constraint in CBD research is limited finance leading to transport problems, inadequate research material, equipment and reduced scientific exchanges. Until recently, there was no pathologist responsible for coffee research in Zimbabwe. The officer carrying this responsibility also had other major crops to research on resulting in inadequate attention being accorded to coffee. There is a need for an equipped mini-laboratory to handle CBD research in the quarantine area to reduce the likelihood of spreading the disease and to alleviate the acute transport problems.

## **NEED FOR REGIONAL AND INTERNATIONAL COOPERATION**

Zimbabwe feels that there is a strong case of regional collaboration as CBD is a common problem in Africa. The collaboration could take the form of germplasm exchange and evaluation, field trials and epidemiological behaviour of the disease in the region. Collaboration may also enable involvement in CBD research by the international community to be spread across a range of beneficiary countries.

## **DISCUSSION**

Responding to participants' reaction to the report, the country's representative, Mr. P. Z. commented that:

- Both coffee berry disease and coffee leaf rust attained economic threshold levels under favourable weather conditions. Use of fungicides was justified as a short term solution to the problem. However, long term control depends on breeding for resistance.
- Trials on growth regulators were aiming at establishing effect of the substances on yield.
- Traditionally, chemical firms in Zimbabwe have been conducting screening and demonstration trials of their pesticides on several crops, coffee inclusive. However, the government does not check the results obtained by the chemical companies.

# **COUNTRY REPORT: MALAWI**

## **COFFEE BERRY DISEASE IN MALAWI**

**NOAH ANTHONY PHIRI**

*Bvumbwe Research Station, P. O. Box 5748, LIMBE, Malawi*

### **INTRODUCTION**

After tobacco, tea and sugar, coffee ranks forth among important export crops in Malawi. Its contribution towards agricultural export earnings is currently estimated at approximately 5.6%.

In Malawi, the total area under coffee is approximately 5,000 hectares, out of which 1,800 ha is under the smallholder subsector mainly in the Northern Region whereas the remaining 3,200 is under estate management, mostly in Mulanje and Thyolo Districts (see Fig 1). Total production of green coffee is about 1,000 MT.

### **COFFEE MANAGEMENT SYSTEMS AND DISEASE LIMITATIONS TO COFFEE YIELD IN MALAWI**

Coffee in Malawi is grown under two main management systems:

- (i) The low input low output management systems which is mainly practised by the smallholder coffee farmers. Under this system, the farmers apply limited amounts of inputs such as pesticides and fertilizers. Under such a management system, diseases are usually endemic.
- (ii) The high input high output management system which is mainly practised in the estate subsector. Under this system, a lot of pesticides are applied resulting in good disease and insect pest control. However, from 1992, the estate subsector has been trying to move away from this kind of management system because of low coffee prices on the world market.

Coffee Leaf Rust (CLR) and Coffee Berry Disease (CBD) are currently the two main diseases of economic importance. CLR has been the major disease in the south where most coffee is grown under the high input high output management system. The major control method has been through the use of preventative and/or curative fungicide sprays. On the other hand, CBD has been the major disease problem in the north. As indicated earlier, inadequate application of fungicides by the smallholder farmers appear to be the main reason for the high disease problem. It is important, therefore, to find cheaper control measures are found for the smallholder sector.

### **COFFEE BERRY DISEASE DISTRIBUTION IN MALAWI**

CBD in Malawi was first confirmed in the Northern Region (Kishyombe and Mainjeni, 1986) in the smallholder coffee sector. Since then, it has spread to most smallholder coffee areas namely, Mulanje Hills, Viphyia North and Nkhatabay Highlands (Fig 1). Until 1991, CBD was confined to smallholder coffee plots in Northern Malawi. The occurrence of the CBD on coffee at Mbanjira Estate, Thyolo District (Fig 1) in 1991 marked the first threat to the coffee industry in Southern region, particularly under estates management. Recently, CBD has also been observed on an estate. Of late, more and more estates have been affected. In a recent survey (Phiri and Mainjeni, unpublished), three more estates have been identified to be affected in the south. The disease is



great to the coffee industry in Malawi. The reduction in inputs by most large coffee farmers brought about by the low world coffee prices is a very unhealthy state as this may lead to unchecked spread of disease. This implies that the CBD is now in both smallholder (North) and estate (South) coffee sectors.




Figure 1: Map of Malawi showing coffee growing areas

## RESEARCH ON COFFEE BERRY DISEASE: DISEASE CONTROL STRATEGIES

The research programme on CBD is fairly young because the disease has not been in the country for a long time.

The main research priorities are (not in order of importance):

(i) **Screening fungicides for efficacy against CBD in Malawi**

This is mainly to test the fungicides recommended in other countries for their effectiveness under Malawi conditions. New fungicides on the market are also included for evaluation. In addition, work on fungicide spray frequencies is also being conducted in order to determine the optimum number of sprays per season.

(ii) **Laboratory and field screening of varieties/lines and individual coffee plants for resistance to CBD**

This area includes work on field selection (selection of individual trees which show resistance) in coffee gardens and their progeny are subjected to preselection tests for confirmation. This work is starting this season - seed collection is in process.

(iii) **Study of epidemiology of *Colletotrichum kahawae* in Malawi**

In this experiment, we intend to look at the development of CBD in relation to the prevailing weather conditions in Malawi. This is in view of the difference in climatic conditions between Malawi and countries where such studies have been conducted before.

(iv) **Breeding for resistance to CBD**

This is one of the research areas where the pathologist, breeder and the agronomist work together. The coffee pathologist will be screening lines, varieties and progeny from coffee gardens while the breeder will actually be breeding the varieties, and the agronomist will be evaluating them for agronomic traits.

Other research activities related to CBD control are:

(i) **Fungicide spray technology**

This research activity is for testing different fungicide spraying equipment. It includes manually operated Knapsack sprayer (commonly used by smallholder farmers), ULV, and motorised (high volume) sprayers. This work is not of high priority.

(ii) **Cultural control**

In this aspect, we intend to monitor the development of CBD in a pruning experiment (including free growth) and population (spacing) experiment. The aim is to identify pruning systems which do not create favourable conditions for the development and spread of CBD.

## ED FOR COLLABORATION IN CONTROL OF COFFEE BERRY DISEASE

laboration in CBD is extremely essential to avoid duplication of work, and for proper use of scarce resources. Collaboration is already effective in Africa such as in research on Root and Tubers, Rhizome, Quarantine analysis and in Red Locust control just to mention a few research areas.

Suggested areas of collaboration could include the following:

- Exchange visits
- Workshops
- Exchange of plant material/technology including reports
- Training - short term particularly for technical staff

## CONCLUSION

To conclude, this report indicates that the occurrence of CBD in Malawi both on the smallholder and estate sub-sectors, pose a new threat to the whole coffee industry in the region. In the short term, mainly due to escalating cost of chemicals and the general reservation on the use of chemicals worldwide, evaluation of fungicides for the control of CBD appears to be the immediate solution to the problem. However, in the long term, the aim should be to develop disease resistant varieties like K711 which was developed in Kenya. Limited work on these lines has just started in Malawi but expertise is not adequate. This is where a joint effort through a research network to combat diseases of economic importance, would be very beneficial to the Coffee Industry in Malawi.

## DISCUSSION

The report was given after the meeting because the representative for Malawi did not attend.

# COUNTRY REPORT: RWANDA

## SYSTEM OF RESEARCH TO CONTROL THE COFFEE BERRY DISEASE (ANTHRACNOSE) IN RWANDA

ANDRÉ SIBOMANA

*Offices des cafés, B. P. 104, Republiaue Rwandaise*

### 1. GENERAL INTRODUCTION

Presently, Rwanda has 117,000,000 coffee plants and 700,000 coffee farmers. This represents an average of 167 coffee plants per farmer. The whole coffee cultivation of the country is practised in private family estates.

The export of coffee contributes for more than 70% of the national export earnings.

The country produces an annual average of 35,000 tonnes of arabica coffee for export and only 600 tonnes of robusta.

Coffee Berry Disease (Anthracnose) has been in Rwanda for a long time and is found in many regions of the country but more significant damages are found in the coastal regions and along the important rivers of the country. These are zones where the humidity is very high especially in the mornings where fog often covers the hills up to about 10 a.m.

The fundamental research on coffee has not experienced much progress at the level of the National Coffee Research Institute (ISAR) due to lack of adequate staff (Researchers and Technicians) and their instability. Similarly, financial and material means have been insufficient. Few experiments have been carried out concerning screening tests, fertilization, farming methods and phytoprotection against coffee bugs (*Antestiopsis lineaticolis*). The document outlines the research orientations of the National Pilot Coffee Research Plan (1990 - 2000).

The Coffee Board started a massive campaign in 1989 to fight this disease through the use of copper oxychloride. Also a research-development programme was undertaken to test different fungicides capable of eradicating this disease. This concerns mainly oxychloride, Dacobre, Topain, Kocide and Daconil.

### 2. NATIONAL RESEARCH PROGRAMME ON COFFEE IN GENERAL AND ON ANTHRACNOSE IN PARTICULAR

#### 2.1 Constraints

##### 2.1.1 Pests and Diseases

- the coffee bug is the most destructive pest (*Antestiopsis lineaticolis*)
- leaf rust (*Hemileia vastatrix*)
- anthracnose (*Colletotrichum kahawae*)
- die-back on old plantations on poor soils

## **2 Farming Practices**

- lack of mulch materials
- bad pruning
- poor soil fertility
- very poor crop density especially in aged plantations

## **3 Post-harvest Technology**

- Bad conditions of pulp removal and drying which have a high influence on the quality of coffee for export.

## **Objectives of the Research**

Since coffee represents the main export crop of the country, efforts in research and extension work must always be undertaken to find solutions to the different production constraints. For the period 1990 - 2000 research must be centred on the following objectives:

- getting early high-yielding varieties;
- obtaining resistant varieties or tolerant to the major pests and disease;
- acquisition of varieties adapted to the different coffee producing areas;
- obtaining varieties whose organoleptic characteristics comply with world market norms;
- implementation of farming methods adapted to agro-ecological conditions of the main production areas: density, pruning, soil shading, fertilization and other appropriate cultivation systems;
- the use of methods to control the main coffee pests and diseases;
- obtaining post-harvest technologies to improve the quality coffee.

## **Research Activities**

### **1 Screening and varietal improvement**

#### **Germplasm**

- Maintenance and enrichment of the collection at the Institute of Agricultural Sciences in Rwanda (ISAR).

#### **Screening**

- Screening of resistant or tolerant varieties to rust and anthracnose.
- Screening of early high-yielding varieties adapted to large producing areas.

### 2.3.2 Farming Methods

- research on the type of mulch and the optimal mode of application with respect to producing areas and taking into account the local availability of soil covering material;
- carrying on with research on living mulch;
- carrying on with research on a better population density according to the varieties (normal habit, reduced habit) and the agro-ecological production zones;
- carrying out with research for the increase of the density of old coffee plants by interplanting young coffee plants;
- comparing the pruning techniques used in Rwanda and elsewhere (East Africa);
- making a synthesis of available data on coffee plants fertilization in the country and evaluating the yield on production;
- research on methods adapted to the regeneration of old coffee plants

### 2.3.3 Crop Protection

- epidemiological studies;
- determination of the effect of incidence on production;
- availability of tolerant or resistant varieties
- experimenting new control and prevention possibilities and evaluation of their effect on yield

### 2.3.4 Post Harvest Technology of Coffee

- Improvement of post-harvest techniques and assessment of their economic impact.

### 2.3.5 Production of Basic Seeds

- Dissemination of new varieties by OCIR

## 3. RESEARCH STAFF IN COFFEE PROGRAMME IN GENERAL

- Researchers: No specific researchers are attached to the research on coffee anthracnose.
- Presently, two researchers work in the programme. Two additional researchers would be assigned to the programme as scheduled.
- Specialists in the following fields would be called in on full or part-time basis:
  - Phytotechnology
  - Screening
  - Entomology
  - Phytopathology
  - Technology

- Part-time researchers are available for entomology and phytopathology.
- **Technicians:** Presently the coffee programme has two technicians.
- The number of technicians would double that of permanent researchers making 8 technicians.

## INFRASTRUCTURES

The Research Institute (ISAR) has a certain number of well-equipped laboratories to carry out its functions conveniently. These are support units to the different programmes including that of coffee. This includes mainly the laboratories of: General Agricultural chemistry, Soil Chemistry and Physics, Phytoprotection, Microbiology, Food Technology, *in vitro* Cultivation, Biometry, Climatology Service and Documentation Service.

ISAR has enough land on its stations to carry out tests.

The programme is financed by the Government of Rwanda and the French Development Fund.

## RESEARCH STATIONS

ISAR has four stations (Rubona, Tamira, Rwere and Karama) but the activities of the coffee programme are carried out only at Rubona South of the country. Meanwhile, tests on the different themes are carried out also in other sites of the country namely: Mayaga (South), Banks of Lake Kivu (West of the country), Mutara (North-East) and Bugesera (Centre-South).

## AVAILABILITY OF DOCUMENTATION

Research documents are kept at the Rubona centre and at the library of the National University, Rwanda. Furthermore, the Institute (ISAR) subscribes to international reviews and periodicals on coffee research.

## SCIENTIFIC EXCHANGES

These exchanges are fairly intensive: training, refresher courses and seminars in different countries especially the USA, France (IRCC), Kenya, Portugal. Exchange of plant materials is also advantageous and ISAR has in its collection, plant material from Portugal, Burundi, Brazil, Ethiopia and Kenya.

Co-operation is done with IRAZ and National Research Institutes of the CEPGL countries.

## TRANSFER OF TECHNOLOGIES AND THEIR IMPACT ON THE FIELD

The transfer of technologies is ensured by extension human resources of the Ministry of Agriculture: transfer of senior staff to operational units assisted by specialized technicians in coffee (belonging to the Coffee Board).

The general assignment of extension senior staff is as follows:

- Identification of regional potentials and constraints in agricultural development.
- Drawing up sub-sectorial policies compatible with regional regulations.
- Establishing realistic regional agricultural development programmes.
- Management of communal services.

In addition to these planning and programming activities of regional agricultural extension service staff are responsible for the following technical activities:

- Organizing research-development activities (analysis of farming techniques as well as study of motivating farmers technically and economically);
- Undertaking the training of communal staff;
- Supervising activities.

Responsibilities during the process of development and transfer of technologies follows:

**a) Diagnostic phase**

This phase concerns collaboration between the Research Institute, extension structure and farmers. This diagnostic is integrated within a precise framework which must be from the onset. Once farming exploitation systems and their constraints are known, it is possible to formulate solutions to these constraints. Some solutions shall be found from new developments already in use and only the verification of adaptation factors suffice. Other constraints will be dealt with through specific research in research stations.

**b) Tests in stations**

These tests, given the scientific rigour required to carry them out, must be under the responsibility of research.

**c) Multilocal tests**

These tests are carried out to take account of the different environmental conditions to ensure the success of innovations to disseminate. In order to attain this objective, tests are conducted at the local level, on permanent multilocal sites or with farmers for this purpose.

Presently, these tests are under research but proposals have been made to improve them by making the extension resources take part in the tests which shall be established at the level of official units.

**d) Adaptation tests**

These are control tests of acceptability which are carried out to verify the pertinence of an innovation according to the conditions of the different exploitation technologies, on a small scale with a rigorous but simple protocol, which must be drawn up jointly with researchers.

The first responsibility of these tests come under the extension structure.

**e) Demonstration and follow-up of themes**

This phase is done through a demonstration training activity centred on identified themes. It is the responsibility of the extension structure. It will enable one to adopt the recommended solutions.



For this last case, a feed-back of research is indispensable, hence it is equally necessary that the protocol of demonstrations be submitted for research approval also.

## CONSTRAINTS

- The research staff is still insufficient and unstable. It suffices to undertake a career in research.
- All researchers must benefit from the 3rd cycle university training.
- The financing of research is not consistent. The coffee sector must support the research programmes relating to coffee in order to ease the granting of all necessary material means.

## DISCUSSIONS

A. Sibomana (*Rwanda*). In Africa there are many regional organizations for instance in March 1993, a network on coffee was established in London by IACO. Then we have the OAU. How are the two going to work together?

N. Nkouka (*OAU*). We shall look at this observation under collaboration.

D. M. Masaba (*Kenya*). I have heard of this cultivar called Jackson five. How is its spread in Rwanda?

A. Sibomana. Jackson two and three other cultivars are widely grown but Jackson five is not yet recommended to the farmers. One of the four cultivars is highly susceptible to CBD.

C. O. Agwanda (*Kenya*). Some of the cultivars in Rwanda are highly susceptible to CBD. Are you trying any other coffee materials from Kenya, Tanzania, etc?

A. Sibomana. Yes, but still under evaluation.

# COUNTRY REPORT: CAMEROON

## PRESENTATION ON ARABICA COFFEE BERRY DISEASE (ANTHRACNOSE) IN CAMEROON

J. BAKALA

IRA/CRA de Nkolbisson, B. P. 2067, Yaounde, Cameroon

### GENERAL SITUATION

Arabica coffee is cultivated in family plantations in Cameroon in altitudes ranging from 1,200 m in the high plateaux of the West and North-West Provinces. This crop covers an area of 16 with a yield of about 20,000 tonnes of coffee every year for export.

The Coffee Berry Disease (Anthracnose) caused by *Colletotrichum kahawae* Waller & Bridg. is the most serious disease which attacks this crop.

Known in Cameroon since 1958, this disease is very serious in areas above 1,500 m in altitude with a favourable climate, it destroys about 60,000 ha of coffee plants and causes a loss in production of 80%.

This loss is so high because the plantations in high altitudes are almost exclusively made up of the Jamaican variety which is very susceptible to this disease.

Damages which the disease causes in lower altitude (1,200 - 1,500 m) regions are of minor importance because it occurs in an uneven manner being more serious in micro-climate regions with a humid mountainous climate.

Associated food and garden crops now yielding much more money than coffee constitute a major handicap to this crop.

### RESEARCH PROGRAMME ON ARABICA COFFEE

The objective is to improve the quantity and quality of Arabica coffee production. In order to achieve these goals, work has been under way since almost twenty years now in the following areas:

**Genetics - Screening** (Varietal tests, hybrid tests, interspecific hybridization, etc ...)

Screening criteria include: vigour, early maturity, productivity, anthracnose and rust tolerance, resistance and the quality of coffee for export.

#### **Agronomy**

Tests for the improvement of cultural techniques, density tests, pruning tests, soil protection, mulching or cover crops and fertilization.

tosanitary protection

chemical control of antestia, coloborrhis and defoliatory caterpillars. Study of the sensitivity of plant material vis-a-vis anthracnose and rusts, chemical control of these diseases.

Technology-Biochemistry

Study of the "stinky bean". The production of coffee quality maps.

## **ATION - INFRASTRUCTURE, EQUIPMENT AND FINANCING**

Research on anthracnose is carried out in two research stations and one research unit:

The Nkolbisson Agricultural Research Station, situated in Yaounde at about 300 km from the arabica area;

The Foumbot Agricultural Research Station situated in the arabica area at an altitude of 1,200 m;

The Santa Research Unit situated in the arabica area at an altitude of 1,900 m and at about 50 km from the Foumbot Station.

Whereas the Santa Unit harbours only collections and test fields, the Foumbot Station has a field laboratory and basic necessary equipment for a phytopathologist.

The Nkolbisson station has a well equipped laboratory where a variety of research work in phytopathology can be performed.

Up to 1987, the working budget of the Coffee Programme in general, and research on anthracnose in particular, was supported by the Cameroon Government but with the economic crisis now plaguing the country, this budget does not exist any more. However, a lot of work has been shelved awaiting attention except fungicide tests intended for chemical control which are financed from time to time by fungicide firms or pesticide distributors. It is through this irregular financing that few field operations have been undertaken.

## **STAFF WORKING ON ANTHRACNOSE**

We have two phytopathologists, two technicians and about 10 field workers working on anthracnose in different research structures.

## **DOCUMENTATION**

There is a well equipped and easily accessible library at the Nkolbisson Agricultural Research Station which, of recent, was receiving numerous scientific journals from countries where research on arabica coffee is undertaken. But since we started experiencing the economic crisis, only some rare journals are being received. The purchase of most of these periodicals requires funds which we do not have any more.

## **CO-OPERATION - EXCHANGE AND TRAINING**

Concerning research on arabica coffee, the Institute of Agricultural Research maintains solid close co-operation ties with CIRAD - ICP which has always placed expatriate researchers at its disposal to achieve certain research work.

We carry out exchange of plant material with all countries producing arabica coffee in the world.

Besides CIRAD, we cooperate with the university of Yaounde within the framework of the Biotechnology Centre where presently, studies on early anthracnose tests are underway. Two research projects on this same disease are in preparation with the cooperation of ORSTOM Portugal (CIFC, OIERAS Portugal) respectively.

Junior staff in the laboratory and those who carry out field observations receive in-service training.

Our researchers and technicians are trained in national or foreign universities through CIRAD Schools of Agriculture in the country respectively. Refresher and short term training courses at these levels are scarce.

## **DISSEMINATION OF RESEARCH FINDINGS ON ARABICA COFFEE**

Counselling to arabica coffee farmers is ensured by two powerful cooperatives and the traditional extension services of the Ministry of Agriculture with which the researchers maintain close contact through visits and training for the transfer of research findings to the beneficiaries. Phytosanitary chemicals purchased every year are recommended after research tests have been carried out jointly with all the parties concerned.

## **CONSTRAINTS RESEARCH NEEDS ON ANTHRACNOSE - COMMENTS AND RECOMMENDATIONS**

One single constraint has shelved research on anthracnose in Cameroon. This constraint is economical. The sharp drop in the price of coffee and other basic products of our country is responsible for the economic crisis which is hitting us. This economic disorder does not make it possible either to our Government or to financing bodies to invest on research on the basic products in general and on arabica coffee in particular.

The price of coffee in the world market must therefore be increased because this is the one condition which may enable the rehabilitation of the arabica crop in general and look for funding for research on anthracnose which is our major constraint in the production of arabica coffee.

For the past years, the African International Scientific Community is becoming more and more united. That Unity is strength and it is finally necessary to unite and co-operate in order to find solutions to the problems of our agriculture which is the pillar of our economies.

Promising initiatives have been taken and many technical networks have been set up. Unfortunately many of these networks are not existing due to lack of funds to revive them and especially to support their researchers work. This is why we think that as far as the process of Scientific Co-operation in Africa is concerned, we should not only create networks but we should also look for funding agencies to keep them alive.

## DISCUSSIONS

Bakala indicated that farmers preferred organics such as Chlorothalonil over copper based formulations due to less labour requirements by the former during fungicide application in the field. The reduced number of sprays of the organic fungicides relative to copper formulations did not seem to lead to higher disease incidence.

## GENERAL DISCUSSIONS ON ALL REPORTS

The participants noted the need to observe information emanating from FAO/WHO concerning banned pesticides. They also shared a common view that installation of a data base system facilitate easy communication between scientists and subsequently help in the standardization of research techniques on common problems.

**P. Zvoutete (Zimbabwe).** Zimbabwe has strict quarantine measures within the country. No coffee zones with CBD is allowed to move to zones with no CBD record.

**N. Nkouka (OAU).** How many countries have quarantine stations?

- Kenya - Muguga
- Zimbabwe - Harare
- Tanzania - TPRI - Arusha
- Ethiopia - Addis Ababa

**D. M. Masaba (Kenya).** CBD was reported to be more serious along the coastal area in Cameroon. What may be contributing to this?

**J. Bakala (Cameroon).** It is probably the high relative humidity coupled with relatively high temperatures 20 - 24°C.

## **PARTICIPANTS**

### **NAME & ADDRESS**

1. **Mr. Eshetu Derso,**  
Research Officer  
Institute of Agricultural Research Centre, Jimma,  
P. O. Box 2003, Addis Ababa.  
Tel: 112695 Jimma.
2. **Mr. Petros Zvoutete,**  
Research Officer  
Plant Protection Research Institute,  
P. O. Box 8100, Causeway, Harare, Zimbabwe.  
Tel: 704531, Telex 22455 AGRIC ZW
3. **Mr. Sibomana André,**  
Chef Service de la Production  
Offices des Cafes,  
B. P. 104, Republique Rwandaise.  
Tel: 23 31 05, Telex 1140 KN
4. **Mr. Joseph Bakala,**  
Plant Pathologist  
IRA/CRA de Nkolbisson  
B. P. 2067, Yaounde, Cameroon.  
Tel: 23 31 05, Telex 1140 KN
5. **Mr. F. S. Ngulu,**  
Plant Pathologist  
Lyamungu Research Institute,  
P. O. Box 3004, Moshi.  
Tel: 4411, Telex 43000 COBOT
6. **Dr. (Mrs.) Agnes Mwang'ombe,**  
Senior Lecturer  
Department of Crop Science,  
University of Nairobi.  
P. O. Box 30197, Nairobi, Kenya,  
Tel: 632175
7. **Dr. (Mrs.) Dinah M. Masaba,**  
Senior Principal Research Officer  
Coffee Research Foundation,  
P. O. Box 4, Ruiru, Kenya.  
Tel: (0151) 21047/92

8. **Mr. Charles Otieno Agwanda,**  
Plant Breeder  
Coffee Research Foundation,  
P. O. Box 4, Ruiru, Kenya.  
Tel: 21047/92
9. **Mr. Phillip Ngotho King'ori,**  
Senior Research Officer  
Coffee Research Foundation,  
P. O. Box 4, Ruiru, Kenya.  
Tel: 21047/92
10. **Dr. N. Nkouka,**  
Ag. Scientific Secretary,  
OAU/IAPSC,  
P. O. BOX 4170, Yaounde, Cameroon.  
Tel: 22 25 28, Telex 8460 KN, Fax (237) 22 54 56

## **SECRETARIAT**

1. **Miss Madrine Rugano,**  
Coffee Research Foundation,  
P O Box 4, Ruiru, Kenya.
2. **Mrs. Njonje Rose,**  
OAU/IAPSC,  
P. O. Box 4170, Yaounde, Cameroon.
3. **Mr. Ngouanet Etienne,**  
OAU/IAPSC,  
P. O. Box 4170, Yaounde, Cameroon

## **OTHER GUESTS**

1. **Hon. Manduku**  
Assistant Minister  
Minister's Representative  
Ministry of Agriculture, Livestock Development & Marketing (MAL&M).
2. **Mr. N. E. S. Muyonga,**  
Deputy Secretary, MAL&M.
3. **Mr. Pithon Mwangi,**  
Chairman, Coffee Board of Kenya (CBK).



## **PARTICIPANTS**

4. **Dr. Wilson R. Opilé,**  
Director of Research  
Coffee Research Foundation (CRF).
5. **Mr. Ole Murunga,**  
General Manager, CBK.
6. **Mr. M. Njiru,**  
CRF Board Member - representing Chairman, CRF Board.
7. **Mr. S. Kloko,**  
Chairman, Kenya Planters Co-operative Union Ltd (KPCU).
8. **Mr. J. K. Gitau,**  
KPCU.
9. **Mr. J. M. King'ang'i,**  
KPCU.
10. **Mr. B. W. Ngundo,**  
Representative - Kenya Agricultural Research Institute (KARI),
11. **Mr. K. A. Mogol,**  
Secretary, Pest Control and Products Board (PCPB).
12. **Prof. D. M. Mukunya,**  
Dean, Faculty of Agriculture,  
University of Nairobi.
13. **Prof. Ole Mbatha,**  
Department of Economics,  
University of Nairobi.
14. **Mr. Fred Ochwoka,**  
Kenya Broadcasting Corporation (KBC).
15. **Mrs. Gatumbi,**  
Representing Senior Plant Pathologist,  
KARI.
16. **Mr. J. K. Mbatha,**  
Chairman, Pest Chemicals Association of Kenya (PCAK).
17. **Mr. J. K. Kinoti,**  
Field Services Manager, CBK.
18. **Mr. J. Muasya,**  
General Manager, Small Scale Coffee Improvement Programme (SCIP II).

19. **Mr. Jacques Colombani,**  
Directeur de Recherche,  
Reprentat de Orstom pour Afrique de Est.
20. **Miss A. W. Wainaina,**  
Head, Entomology Section,  
CRF.
21. **Mr. M. P. H. Gathaara,**  
Head, Crop Physilogy Section,  
CRF.
22. **Dr. J. Mburu Njoroge,**  
Head, Experimental Agronomy Section,  
CRF.
23. **Mr. A. M. Karanja,**  
Head, Agricultural Economics Section,  
CRF.
24. **Mr. Mburu Ngige,**  
Head, Chemistry Section,  
CRF.
25. **Mr. A. S. K. Maithia,**  
Farm Manager,  
Azania Estate, CRF.
26. **Mr. N. D. Njeru,**  
Deputy Head,  
Research Liaison Section, CRF.

AFRICAN UNION UNION AFRICAINE

African Union Common Repository

<http://archives.au.int>

Organs

African Union Commission

1995-08

# African journal of plant protection

Organization of African Unity

Organization of African Unity

<http://archives.au.int/handle/123456789/6402>

*Downloaded from African Union Common Repository*